

The Inhibition of NF- κ B Activation Pathways and the Induction of Apoptosis by Dithiocarbamates in T Cells Are Blocked by the Glutathione Precursor N-Acetyl-L-Cysteine

Paula C. Fernandez¹, Joel Machado Jr.¹,
Volker T. Heussler¹, Catherine Botteron¹,
Guy H. Palmer² and Dirk A.E. Dobbelaere^{1,*}

¹ Laboratory of Molecular Pathology, Institute of Animal Pathology, University of Berne, Länggass-Str. 122, CH-3012, Berne, Switzerland

² Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, WA 99614-7040, USA

* Corresponding author

Nuclear factor- κ B regulates genes that control immune and inflammatory responses and are involved in the pathogenesis of several diseases, including AIDS and cancer. It has been proposed that reactive oxygen intermediates participate in NF- κ B activation pathways, and compounds with putative antioxidant activity such as N-acetyl-L-cysteine (NAC) and pyrrolidine dithiocarbamate (PDTC) have been used interchangeably to demonstrate this point. We examined their effects, separately and combined, on different stages of the NF- κ B activation pathway, in primary and in transformed T cells. We show that NAC, contrary to its reported role as an NF- κ B inhibitor, can actually enhance rather than inhibit I κ B degradation and, most importantly, show that in all cases NAC exerts a dominant antagonistic effect on PDTC-mediated NF- κ B inhibition. This was observed at the level of I κ B degradation, NF- κ B DNA binding, and HIV-LTR-driven reporter gene expression. NAC also counteracted growth arrest and apoptosis induced by dithiocarbamates. Antagonistic effects were further observed at the level of jun-NH₂-terminal kinase, p38 and ATF-2 activation. Our findings argue against the widely accepted assumption that NAC inhibits all NF- κ B activation pathways and shows that two compounds, previously thought to function through a common inhibitory mechanism, can also have antagonistic effects.

Key words: Antioxidant / Disulfiram / Pyrrolidine dithiocarbamate / Reactive oxygen intermediates / T cell transformation / Theileria.

Introduction

NF- κ B binds to decameric κ B sites which are found in the promoters and enhancers of genes that are involved in in-

flammation, immune responses, disease, and apoptosis (Baeuerle and Henkel, 1994; Siebenlist *et al.*, 1994; Baldwin, 1996). Consequently, extensive efforts are being invested in identifying steps in the NF- κ B activation pathway that could be blocked in a targeted manner (Poli and Fauci, 1992; Blackwell *et al.*, 1996; Sakurada *et al.*, 1996). NF- κ B is activated by a wide array of agents including cytokines, oxidants such as H₂O₂, phorbol esters, certain phosphatase inhibitors such as okadaic acid (OA), and UV or ionising radiation. Different receptors, among them members of the tumour necrosis factor receptor family that participate in apoptosis (Nagata, 1997), also activate NF- κ B. In addition, NF- κ B may participate in the pathogenesis of diseases caused by viruses and bacteria (reviewed in Baeuerle and Henkel, 1994; Baldwin, 1996). More recently, the list of NF- κ B inducers has been extended to parasites such as *Toxoplasma* (Gazzinelli *et al.*, 1996), *Plasmodium* (Schofield *et al.*, 1996) and *Theileria* (Ivanov *et al.*, 1989; Palmer *et al.*, 1997) which interfere with host cell signalling pathways.

The NF- κ B/Rel family of proteins consists of p50 (NF κ B1), p52 (NF κ B2), p65 (RelA), c-Rel and RelB (reviewed in Verma *et al.*, 1995; Baeuerle and Baltimore, 1996; Baldwin, 1996). These proteins share a conserved 'Rel homology domain' through which they can form homodimers or heterodimers which bind DNA. In most cell types, NF- κ B is sequestered in the cytoplasm in a latent form in association with inhibitory I κ B proteins, of which I κ B α and I κ B β are the best characterised. Activation of NF- κ B has been shown to involve signal-induced I κ B phosphorylation by I κ B kinases (reviewed in Stancovski and Baltimore, 1997). I κ B phosphorylation and subsequent ubiquitination targets the protein for proteosomal degradation (Palombella *et al.*, 1994; Traenckner *et al.*, 1994; DiDonato *et al.*, 1996). Ubiquitination-independent I κ B degradation (Lee *et al.*, 1997a) also occurs, however, and may be involved in regulating basal I κ B turnover (Krappmann *et al.*, 1996).

Numerous studies suggest that reactive oxygen species (ROS) serve as common intracellular mediators of I κ B degradation and subsequent NF- κ B activation (Schulze Osthoff *et al.*, 1993; Baeuerle and Henkel, 1994; Baeuerle *et al.*, 1996; Sen and Packer, 1996; Flohé *et al.*, 1997). Consistent with this hypothesis, it has been shown that a variety of structurally diverse antioxidants are capable of inhibiting NF- κ B activation. A number of studies, however, also suggest that not all NF- κ B activation pathways involve ROS, as judged by their resistance to antioxidants (Suzuki *et al.*, 1994; Brennan and O'Neill, 1995;

Flohé *et al.*, 1997). N-acetyl-L-cysteine (NAC) and pyrrolidine dithiocarbamate (PDTC) are two compounds that have been used extensively as NF- κ B inhibitors (Schreck *et al.*, 1992; Meyer *et al.*, 1994). NAC, a glutathione (GSH) precursor and ROS scavenger, inhibits NF- κ B activation induced by a range of stimuli in several, but not all, cell types (Schreck *et al.*, 1991; Meyer *et al.*, 1993). PDTC, a dithiol compound often used because of its reported antioxidant activity, is a more consistent inhibitor of NF- κ B which appears to act more independently of the activating agent and cell line used (Schreck *et al.*, 1991; Galter *et al.*, 1994; Mihm *et al.*, 1995). PDTC has recently also been shown to induce G1 cell-cycle arrest and apoptosis in malignant cell lines and to enhance the anti-tumour efficacy of chemotherapeutic agents, prompting the proposal that PDTC may find application in combination therapy (Chinery *et al.*, 1997).

The fact that NAC and PDTC can independently inhibit NF- κ B activation has been used in support of a general model in which NF- κ B activation is governed by oxidative stress (Schreck *et al.*, 1991). Evidence is emerging, however, that the major mode of inhibitory action of PDTC may not be restricted to its antioxidant activity, but could also include metal ion chelating (Bessho *et al.*, 1994; Brennan and O'Neill, 1995) and even pro-oxidant properties (Nobel *et al.*, 1995; Brennan and O'Neill, 1996b; Pinkus *et al.*, 1996). In this context, PDTC-dependent activation of the transcription factor AP-1 was shown to depend on its pro-oxidant activity and could be blocked by anti-oxidants, including GSH and NAC, its synthetic precursor (Pinkus *et al.*, 1996). In the present study, we investigated whether such antagonistic effects also existed at the level of the NF- κ B activation in primary lymph node (LN) T cells and on induced and constitutive NF- κ B activation in transformed T cells. We report that the GSH precursor, NAC, which is generally accepted as an antioxidant inhibitor of NF- κ B, can actually enhance NF- κ B activation rather than inhibit it, and, more importantly, we demonstrate that the inhibitory effects of dithiol compounds such as PDTC and disulfiram on NF- κ B activation are neutralised completely by NAC. This effect can be observed at the level of I κ B α and I κ B β degradation, NF- κ B DNA binding activity, NF- κ B transcriptional activity and dithiocarbamate-induced growth arrest and apoptosis. Our findings demonstrate that these classical NF- κ B inhibitors, previously thought to function through a common antioxidant mechanism, can have completely diverging and antagonistic effects on NF- κ B activation pathways.

Results

NAC Enhances Basal I κ B Degradation

I κ B α , in particular the uncomplexed form, is inherently unstable and undergoes a continuous turnover in the absence of NF- κ B-inducing signals (Miyamoto *et al.*, 1994; Krappmann *et al.*, 1996). This is most apparent when *de*




Fig. 1 Effect of NAC and PDTC on I κ B Degradation in Resting LN T Cells and Transformed Tpi T Cells.

(A) Basal I κ B α turnover in LN T cells is enhanced by NAC. LN T cells were cultured in the presence or absence of CHX to inhibit *de novo* I κ B α synthesis and treated with NAC or PDTC, as indicated. Whole cell extracts (15 μ g/lane) were examined by immunoblot analysis using anti-I κ B α Ab. The position of I κ B α is indicated by an open triangle.

(B) Effect of NAC and PDTC on constitutive I κ B degradation in Tpi T cells. Tpi T cells were cultured in the presence of CHX, NAC or PDTC as indicated and whole cell extracts (20 μ g/lane) examined by immunoblot analysis using anti-I κ B α and anti-I κ B β Ab together. The positions of I κ B α and I κ B β are indicated by open triangle and closed triangles, respectively. The upper open triangle marks phosphorylated I κ B α .

novo synthesis is blocked by the protein synthesis inhibitor cycloheximide (CHX). When resting LN T cells were cultured in the presence of CHX (Figure 1A), basal levels of I κ B α turnover could be demonstrated by a modest reduction in the amounts of I κ B α protein detected by Western blot analysis. Whereas PDTC treatment showed no effect, NAC actually enhanced I κ B α turnover, indicating that the two compounds affect basal I κ B α turnover differently. This finding was surprising, since NAC, as an antioxidant and ROS scavenger, was expected to inhibit rather than enhance I κ B α degradation.

NAC Blocks PDTC-Mediated Inhibition of I κ B Degradation

In order to investigate whether this observation could be confirmed in cells in which the NF- κ B activation pathway is constitutively activated, we next tested the effect of NAC and PDTC on I κ B degradation in transformed T cells. As a source of transformed T cells, we used *Theileria parva*-infected T cells (Tpi T cells) which proliferate continuously and behave as persistently activated T cells (Dobbelare *et al.*, 1988; Eichhorn *et al.*, 1993; Galley *et al.*, 1997). In Tpi T cells, NF- κ B is constitutively activated, involving the continuous degradation of both I κ B α and I κ B β and nuclear translocation of several members of the NF- κ B/Rel family. Steady-state levels of I κ B were not sig-

nificantly affected by NAC or PDTC treatment except for a slight reduction in the amount of phosphorylated $\text{I}\kappa\text{B}\alpha$ in PDTC-treated cells (data not shown). Blocking *de novo* protein synthesis in Tpi T cells by treatment with CHX for 4 h resulted in decreased levels of both $\text{I}\kappa\text{B}\alpha$ and $\text{I}\kappa\text{B}\beta$ compared to untreated cells (Figure 1B), reflecting the constitutive degradation of $\text{I}\kappa\text{B}$ (Palmer *et al.*, 1997). As was observed for basal degradation in primary LN T cells, NAC enhanced constitutive $\text{I}\kappa\text{B}\alpha$ degradation. Treatment with PDTC clearly prevented $\text{I}\kappa\text{B}$ degradation. Furthermore, in the presence of NAC, the inhibition of $\text{I}\kappa\text{B}$ degradation by PDTC was totally abrogated, demonstrating that the antioxidant NAC can block the effect of PDTC at the level of $\text{I}\kappa\text{B}$ degradation. OA is a strong activator of NF- κB and induces the rapid phosphorylation and degradation of $\text{I}\kappa\text{B}$ (Rieckmann *et al.*, 1992; Menon *et al.*, 1995), most likely by blocking protein phosphatase 2A which inactivates $\text{I}\kappa\text{B}$ kinase (DiDonato *et al.*, 1997). In earlier work, it has been proposed that OA induces $\text{I}\kappa\text{B}$ degradation downstream of the anti-oxidant-sensitive step in the NF- κB activation cascade (Suzuki *et al.*, 1994; Sen and Packer, 1996). When resting LN T cells were stimulated with OA, $\text{I}\kappa\text{B}\alpha$ was phosphorylated and rapidly degraded (Figure 2A) and in agreement with earlier reports (Suzuki *et al.*, 1994; Sen and Packer, 1996), $\text{I}\kappa\text{B}\alpha$ degradation was not inhibited by NAC, PDTC or a combination of both. OA-induced $\text{I}\kappa\text{B}$ degradation was also restricted to $\text{I}\kappa\text{B}\alpha$ and no $\text{I}\kappa\text{B}\beta$ degradation could be detected. In the transformed Tpi T cells, OA also induced the complete degradation of




Fig. 2 Effect of NAC and PDTC on OA-Induced $\text{I}\kappa\text{B}$ Degradation in Resting LN T Cells and Transformed Tpi T Cells.

(A) NAC and PDTC have no effect on OA-induced $\text{I}\kappa\text{B}\alpha$ degradation in resting LN T cells. LN T cells were stimulated with OA in the presence or absence NAC or PDTC as indicated. The position of $\text{I}\kappa\text{B}\alpha$ (lower open triangle) and its phosphorylated form (upper open triangle) and the position of $\text{I}\kappa\text{B}\beta$ (closed triangle) are indicated.

(B) Tpi T cells were stimulated with OA in the presence or absence of NAC and PDTC and protein extracts were analysed for $\text{I}\kappa\text{B}$ levels as above.

$\text{I}\kappa\text{B}\alpha$ (Figure 2B). In contrast to resting LN T cells in which $\text{I}\kappa\text{B}\beta$ levels were not affected by OA, significant $\text{I}\kappa\text{B}\beta$ degradation could be detected in transformed Tpi T cells. Only a faint $\text{I}\kappa\text{B}\beta$ band (small closed triangle) could be detected upon OA treatment, and degradation was again enhanced by NAC. PDTC largely prevented the OA-induced degradation of $\text{I}\kappa\text{B}\beta$, but not that of $\text{I}\kappa\text{B}\alpha$. The stabilisation of $\text{I}\kappa\text{B}\beta$ conferred by PDTC was completely prevented by NAC, demonstrating that the antagonistic effect of NAC on PDTC can occur both at the level of constitutive $\text{I}\kappa\text{B}$ turnover and OA-induced $\text{I}\kappa\text{B}$ degradation. Triggering of T cell surface receptors for TNF, $\text{IFN}\gamma$, $\text{IL1}\beta$ or stimulation of CD95/Fas or CD28 generates ceramide via hydrolysis of sphingomyelin by sphingomyelinases (reviewed in Ballou *et al.*, 1996). Ceramide is an important signal transducer of NF- κB activation. N-acetylsphingosine, a membrane-permeable synthetic form of ceramide, was used to test the effect of PDTC and NAC on ceramide-induced $\text{I}\kappa\text{B}$ degradation in Tpi T cells. Within 1 h of ceramide treatment, $\text{I}\kappa\text{B}\alpha$ was strongly reduced and $\text{I}\kappa\text{B}\beta$ had all but disappeared. PDTC inhibited the ceramide-induced degradation of both $\text{I}\kappa\text{B}\alpha$ and $\text{I}\kappa\text{B}\beta$. NAC further enhanced $\text{I}\kappa\text{B}$ degradation and the PDTC-mediated inhibition of $\text{I}\kappa\text{B}$ degradation was again prevented by NAC (not shown).

The Antagonistic Effect of NAC on PDTC Is Also Evident at the Level of NF- κB DNA Binding and Transcriptional Activity

In previous work we have shown by electrophoretic mobility shift assay (EMSA) that *T. parva* induces NF- κB activation resulting in the appearance of different κB binding complexes in the nucleus (Ivanov *et al.*, 1989). NF- κB transcriptional activity in Tpi T cells was also shown to be resistant to NAC and only partly reduced by treatment with PDTC (Palmer *et al.*, 1997). Consistent with these observations, NAC had no inhibitory effect on NF- κB DNA binding activity (Figure 3); on the contrary, in agreement with the enhanced $\text{I}\kappa\text{B}$ degradation induced by NAC, DNA binding was often moderately increased in the presence of




Fig. 3 Influence of NAC and PDTC on NF- κB DNA Binding Activity.

NF- κB translocation and DNA binding in Tpi T cells is partially inhibited by PDTC, but not by NAC, and NAC counteracts PDTC-mediated inhibition. Tpi T cells were left untreated or treated for 5 h with PDTC and NAC as indicated. Nuclear lysates were prepared from 10^7 cells and NF- κB binding to [^{32}P]-labelled oligonucleotides representing consensus κB recognition sites determined by EMSA. The arrowhead indicates the NF- κB complex which is most affected by PDTC. The lane labelled 'comp.', shows an assay in which an excess of unlabelled oligonucleotide included in the EMSA as a control for binding specificity.

NAC. PDTC treatment resulted in a partial inhibition of DNA binding, affecting predominantly the faster migrating NF- κ B complex (marked by an arrowhead). NAC completely restored NF- κ B DNA binding in PDTC-treated cells as shown by the reappearance of the lower complex. Treatment of Tpi cells with OA resulted in increased NF- κ B binding activity (not shown) and consistent with the observations made for I κ B, NAC did not inhibit either quantitative or qualitative changes. Treatment with PDTC, on the other hand, resulted in reduced NF- κ B DNA binding activity, but this effect was at least partly blocked when cells were simultaneously treated with NAC (data not shown).

To test whether the antagonistic effects of NAC on the activity of PDTC could also be demonstrated at the level of NF- κ B transcriptional activity, transient transfection experiments were performed using the reporter plasmid HIV-CAT. In this plasmid, expression of the chloramphenicol

acetyl transferase (CAT) gene is placed under the control of the two κ B binding sites within the HIV-LTR. As shown before for Tpi T cells (Palmer *et al.*, 1997), the constitutive NF- κ B transcriptional activity, monitored by CAT activity, was not inhibited by NAC (Figure 4A), but was reduced by 62% when cells were treated with PDTC. No reduction in CAT activity could be observed, however, when cells were treated simultaneously with both compounds. The effects of both compounds on basal NF- κ B transcriptional activity in Jurkat T cells was identical to that observed in Tpi T cells, with CAT activity reduced by 57% in the presence of PDTC (data not shown). In Tpi T cells, NF- κ B is maximally induced and can not be activated further. Stimulation of Jurkat T cells with TPA and ionomycin, however, resulted in a 4.8-fold induction above basal activity (Figure 4B), which was completely inhibited by PDTC. NAC did not interfere with the induction of




Fig. 4 PDTC-Mediated Inhibition of NF- κ B Dependent Reporter Gene Expression Is Abolished by NAC.

(A) Tpi T cells, in which NF- κ B is constitutively activated, were transfected with the plasmid HIV-CAT, in which CAT expression is stimulated through two κ B binding sites. Transfected cells were either untreated or incubated with PDTC and/or NAC for 24 h prior to determination of CAT activity. Results represent one of at least three separate experiments and are expressed as % of CAT activity in untreated HIV-CAT-transfected cells. Bars represent the mean value (\pm S.D., $n=3$).

(B) Effect of NAC and PDTC on NF- κ B-dependent reporter gene expression in HIV-CAT-transfected Jurkat T cells, induced by treatment with TPA and ionophore. Results from one of at least three experiments are presented as fold induction compared to unstimulated Jurkat T cells. Bars represent the mean value (\pm S.D., $n=3$).




Fig. 5 NAC Prevents PDTC-Induced Growth Arrest and Apoptosis.

(A) Jurkat T cells were cultured overnight in the presence or absence of PDTC and NAC at the indicated doses. Cells were labelled with [methyl-³H]thymidine [³H]TdR the last 5 h of culture and [³H]TdR incorporation measured. Data are presented as cpm $\times 10^{-3}$ per 4×10^4 cells seeded. Error bars represent 1 SD of the mean of values from triplicate cultures. (B) Jurkat T cells were cultured for 16 h in the presence of PDTC (10 μ g/ml) alone (solid line) or PDTC and NAC (30 mM) together (dotted line), and analysed by flow cytometry for the binding of fluoresceine-labelled annexin-V. The grey profile shows annexin-V binding to control cells treated with NAC alone. Propidium iodide DNA staining was used to exclude necrotic cells from the analysis.

NF- κ B and again prevented PDTC-mediated inhibition of NF- κ B activation. These data confirm that the antagonistic effect of NAC on PDTC-mediated inhibition of NF- κ B activation can be observed at several levels of the activation cascade.

NAC Counters PDTC-Induced Growth Arrest and Apoptosis in Transformed T Cells

There are conflicting reports on the effect of PDTC on proliferation and apoptosis. PDTC has been reported to induce apoptosis in thymocytes (Nobel *et al.*, 1995) and tumour cells (Chinery *et al.*, 1997; Giri and Aggarwal, 1998). On the other hand, PDTC has also been shown to protect cells from apoptosis (Albrecht *et al.*, 1994; Bessho *et al.*, 1994; Mihm *et al.*, 1995). The proliferation of both Tpi and Jurkat T cells was monitored by [methyl- 3 H]thymidine incorporation and found to be potently inhibited by micromolar amounts of PDTC. In both cases, however, the PDTC-induced proliferation arrest was reversed by cotreatment with NAC (Figure 5A). In order to determine whether growth arrest was followed by apoptosis, the expression of phosphatidylserine was also monitored. During the induction of apoptosis, changes occur at the cell surface, including the translocation of phosphatidylserine (PS) from the inner part of the plasma membrane to the outer layer. Annexin-V is a PS-binding protein which binds with high affinity to PS and can be detected by flow cytometry (Verhoven *et al.*, 1995). When Jurkat T cells were treated for 16 h with PDTC, the majority of cells exposed PS on their surface (Figure 5B). A clear reduction in annexin-V binding could be observed, however, when cells were cultured in the presence of both PDTC and NAC, demonstrating that NAC is capable of preventing PDTC-induced apoptosis in Jurkat T cells. Identical results were obtained for Tpi T cells (not shown).

Effects of NAC and PDTC on jun-NH₂-Terminal Kinase (JNK), p38 and ATF-2

Many stimuli that induce NF- κ B also activate the MAP kinase family members JNK and p38. Previous work has shown that NAC and PDTC can both induce JNK in Jurkat T cells (Gomez del Arco *et al.*, 1996). We tested whether this also applied to resting LN T cells and whether NAC could block PDTC-induced effects. Basal JNK and p38 activity and their induction by osmotic shock were examined. Nuclear extracts from LN T cells were subjected to Western blot analysis, using antibodies (Ab) directed against phosphorylated forms of c-Jun and p38. Phosphorylation of c-Jun, detectable by anti-phospho-c-Jun Ab, reflects JNK activity; likewise, p38 activation is accompanied by specific p38 autophosphorylation. PDTC treatment resulted in increased c-Jun phosphorylation in resting LN T cells and also in cells that were subjected to mild osmotic shock (Figure 6A, lanes 3 and 7). NAC weakly super-induced JNK activation only in NaCl-treated cells. PDTC-mediated JNK activation, however, was potently inhibited when NAC was present (compare lanes 3 and 4,




Fig. 6 Effects of NAC and PDTC on JNK, p38 and ATF-2 Activation.

(A) Effect in resting LN T cells and LN T cells subjected to osmotic shock. Resting LN T cells (lanes 1–4) or LN T cells subjected to osmotic shock (0.2 M NaCl for 20 min; lanes 5–8) were cultured in the presence or absence of PDTC and NAC. Nuclear extracts were analysed by immunoblotting, using Ab specific for phospho-c-Jun (top panel) or phospho-p38 (lower panel). Closed arrows point at the specific phospho-c-Jun and phospho-p38 bands. In the lower panel, the position of a non-specific band (ns) is indicated by an open arrow.

(B) Effect of NAC and PDTC on ATF-2 activation in Tpi T cells subjected to osmotic shock. Immunoblot analysis of extracts prepared from Tpi T cells that were either untreated or submitted to osmotic shock (even numbers) in the presence or absence of NAC and PDTC. The filter was probed with an anti-ATF-2 Ab; unphosphorylated and phosphorylated ATF-2 are indicated by open and closed arrowheads, respectively.

lanes 7 and 8). In Tpi T cells, in which JNK is constitutively activated (Galley *et al.*, 1997), superinduction of JNK by PDTC was also blocked by NAC (data not shown). A basal level of p38 activity could be detected in resting LN T cells which was not observed in the presence of NAC. PDTC treatment did not result in additional p38 activity. In LN T cells subjected to osmotic shock, p38 was clearly upregulated and this was enhanced by NAC or PDTC treatment. In the presence of both compounds, however, the levels of phospho-p38 were markedly reduced.

The JNK and p38 pathways have been reported to regulate the phosphorylation and activation of transcription factors such as AP-1 and ATF-2 (Gupta *et al.*, 1995) (reviewed in Whitmarsh and Davis, 1996). It has previously been described that PDTC-induced AP-1 activation can be inhibited by glutathione and NAC (Pinkus *et al.*, 1996). We investigated whether this also applied to ATF-2. Activation of ATF-2 is accompanied by changes in its phosphorylation pattern, resulting in the appearance of additional bands with reduced electrophoretic mobility. Total cellular extracts from Tpi and resting LN T cells were

subjected to immunoblot analysis, using Ab specific for ATF-2. Marked changes in ATF-2 phosphorylation patterns could be observed in Tpi or LN T cells that were subjected to osmotic shock (shown for Tpi T cells in Figure 6B, even numbers). NAC treatment did not induce any changes, but ATF-2 phosphorylation was more pronounced in the presence of PDTC (Figure 6B, compare lanes 2 and 4) and the pattern of intensity of the different bands was also altered. In the presence of both NAC and PDTC, however, ATF-2 phosphorylation induced by osmotic shock was significantly reduced (Figure 6B, compare lanes 4 and 8). Thus, PDTC can enhance both AP-1 (Pinkus *et al.*, 1996) and ATF-2 activation and, in both cases, these effects are reversed by NAC.

NAC Also Counteracts the Inhibitory Effects Induced by Another Thiol, Disulfiram

Finally, we determined whether NAC could counteract the action of dithiol compounds other than PDTC. Disulfiram, a drug used for alcohol aversion therapy, is also a potent inhibitor of NF- κ B (Schreck *et al.*, 1992). Disulfiram treatment of Tpi T cells resulted in a strong inhibition of NF- κ B DNA binding (Figure 7A). Although the pattern of inhibition was different from that caused by PDTC, disulfiram-mediated

NF- κ B inhibition was completely prevented by NAC. Inhibition of NF- κ B activation by non-thiol compounds such as *o*-phenanthroline and desferrioxamine, on the other hand, was not blocked by NAC (data not shown). Transfection studies with HIV-CAT plasmid showed that disulfiram treatment resulted in a significant reduction in CAT activity (Figure 7B) which was reversed by NAC. As was the case for PDTC, disulfiram caused rapid growth arrest and apoptosis of Jurkat and Tpi T cells, which was counteracted when NAC was added to the culture medium. Figure 7C shows a typical experiment using Tpi T cells in which proliferation was monitored by measuring [methyl- 3 H]thymidine incorporation. Disulfiram potently blocked proliferation, but this did not occur in the presence of NAC. An identical pattern was again observed for Jurkat T cells (not shown).

Discussion

Lack of Susceptibility to Inhibition by NAC

Many inducers of NF- κ B activation are known to result in the generation of intracellular ROS (Schreck *et al.*, 1992; Meyer *et al.*, 1994) (reviewed in Schreck and Baeuerle,




Fig. 7 NAC Blocks Disulfiram-Mediated Inhibition of NF- κ B and Disulfiram-Induced Growth Arrest.

(A) NAC prevents disulfiram-mediated inhibition of NF- κ B activation. Tpi T cells were left untreated or treated with disulfiram (1 μ M) and/or NAC (30 mM) for five hours. Nuclear extracts were examined by EMSA as described for Figure 3.

(B) The inhibition of NF- κ B transcriptional activity by disulfiram is prevented by NAC. Tpi T cells were transfected with the plasmid HIV-CAT. Constitutive NF- κ B transcriptional activity was monitored in transfected cells that were either left untreated or incubated with disulfiram (1 μ M) and/or NAC (30 mM) for 24 hours prior to determining of CAT activity. One of two separate experiments is shown and results are expressed as % of CAT activity in untreated control cells. Bars represent the mean value (\pm S.D., $n = 3$).

(C) Disulfiram-induced arrest of proliferation is prevented by NAC. Tpi T cells were cultured for 24 h in the presence or absence of disulfiram and NAC at the indicated doses. Cells were labelled with [methyl- 3 H]thymidine ([3 H]TdR) for the last 5 h of culture and incorporation measured by liquid scintillation counting. Data are presented as cpm $\times 10^{-3}$ per 4×10^4 cells seeded. Error bars represent 1 SD of the mean of values from triplicate cultures.

1994). This, together with the fact that NF- κ B activation can be inhibited by antioxidants, has led to the proposal that ROS serve as second messengers in most, if not all, NF- κ B activation pathways (Schreck *et al.*, 1991; Schreck and Baeuerle, 1994). In these studies, NAC and dithiols have been used interchangeably to underpin this theory. Thus, both PDTC and NAC have been shown to inhibit the synthesis of IL-6, IL-8, and GM-CSF induced by TNF or bacterial lipopolysaccharides (LPS) in human umbilical vein endothelial cells (Munoz *et al.*, 1996) and similar observations were made in other systems (Lee *et al.*, 1997b; Kikumori *et al.*, 1998). Contrary to our findings, NAC has also been shown to inhibit NF- κ B activation in a number of T cell lines (Dröge *et al.*, 1991; Mihm *et al.*, 1995). The concept that NF- κ B is a sensor for oxidative stress has recently been challenged more directly (Li and Karin, 1999) and arguments against a general involvement of ROS in NF- κ B activation are accumulating (Suzuki *et al.*, 1994, 1995) (reviewed in Flohé *et al.*, 1997; Li and Karin, 1999). NAC failed to inhibit the activation of NF- κ B by classical NF- κ B-inducing agents such as IL-1, TNF and TPA in some cell lines, but not in others (Brennan and O'Neill, 1995; Bowie *et al.*, 1996). Likewise, in the T lymphocytic cell line, CEM, NF- κ B activation by PMA was resistant to NAC (Legrand Poels *et al.*, 1997). These findings indicate that the ROS model of NF- κ B activation and sensitivity to NAC is restricted to certain cell types. Furthermore, H₂O₂ does not invariably activate NF- κ B (Israel *et al.*, 1992; Anderson *et al.*, 1994; Brennan and O'Neill, 1995) and studies involving the transient overexpression of catalase suggest that H₂O₂ may not always serve as a second messenger in NF- κ B activation (Suzuki *et al.*, 1995). Defining the exact role of H₂O₂ (or other ROS) as an integratory secondary messenger for different T cell signals has indeed been complicated by the fact that various T cell lines and peripheral blood T cells can differ markedly – also within one cell line – in the levels of NF- κ B activation induced by oxidant stress (Ginn Pease and Whisler, 1998). Various factors appear to regulate the responsiveness, since it has been shown that optimal NF- κ B-mediated transcriptional responses in Jurkat T cells exposed to oxidative stress are dependent on intracellular glutathione and costimulatory signals (Ginn Pease and Whisler, 1996). The nature of the stimulus may also play a role. For instance, it has been shown that NAC and PDTC were also not able to block daunomycin-induced NF- κ B activation or p53 induction, indicating that ROI are not involved in the cellular response to daunomycin stimulation (Hellin *et al.*, 1998).

A consistent observation in our work was the general lack of inhibitory activity observed for NAC in the systems that were tested. Assuming that NAC is an antioxidant which indeed blocks the effects of ROS, our observations strengthen the concept that the ROS-dependence of NF- κ B activation may not be as strict or widespread as previously proposed (Schulze Osthoff *et al.*, 1997) and also argue in favour of cell-type restriction. Moreover, and contrary to expectations, NAC enhanced rather than inhibited I κ B degradation. Indeed, we found the basal I κ B α turn-

over in resting LN T cells, the constitutive degradation in transformed Tpi T cells, and OA- or ceramide-induced I κ B degradation to be enhanced in the presence of NAC. While this observation may be unexpected, NAC-mediated NF- κ B activation is not without precedent. It has recently been reported that treatment of macrophages with NAC resulted in increased NF- κ B activity which correlated with an up-regulation of HIV production (Nottet *et al.*, 1997).

NAC Blocks PDTC-Mediated Inhibition of NF- κ B

One of the salient observations of our work was that NAC consistently blocked the PDTC-mediated inhibition of the NF- κ B activation pathway, irrespective of the way in which NF- κ B activation was induced. This could be observed at the level of I κ B degradation, NF- κ B DNA binding and transcriptional activity. These findings are not compatible with the proposed common mode of action for PDTC and NAC, in which both compounds are thought to inhibit NF- κ B activation through their antioxidant activity (Meyer *et al.*, 1993, 1994). Does PDTC really function as an antioxidant? Dithiocarbamates can induce a variety of other biological effects in mammalian cells (Sen and Packer, 1996; Flohé *et al.*, 1997). A number of studies suggest that the inhibition of NF- κ B by PDTC and other dithiocarbamates occurs through their metal ion chelating properties (Bessho *et al.*, 1994; Brennan and O'Neill, 1995; Nobel *et al.*, 1995) and several other, non-thiol metal chelators such as *o*-phenanthroline and desferrioxamine also inhibit NF- κ B (Schreck *et al.*, 1992). Nevertheless, we found that the inhibitory effect of the latter two compounds was not blocked by NAC, suggesting that NAC probably does not act by blocking the potential metal chelating properties of PDTC. In addition, it has been proposed that PDTC may cause GSH oxidation, resulting in GSSG accumulation, by a non-radical mechanism (Ziegler, 1991). Brennan and O'Neill (1996a) also reported PDTC pro-oxidant activity, resulting in increased levels of GSSG which can affect the binding of NF- κ B to DNA. In addition, PDTC pro-oxidant, rather than antioxidant, activity has been shown to be responsible for inducing AP-1 activation (Pinkus *et al.*, 1996). NF- κ B activation apparently requires optimal levels of GSSG (Dröge *et al.*, 1994) which is regulated in a complex manner. At low GSSG levels, T cells cannot optimally activate NF- κ B, whereas high GSSG levels inhibit the DNA binding activity of NF- κ B. The effects of GSSG, in turn, are antagonized by reduced thioredoxin (Dröge *et al.*, 1994) which has distinct roles in the cytoplasm and in the nucleus (Flohé *et al.*, 1997). Considered together, it is conceivable that NAC, as a precursor of GSH, counteracts the effects of increased GSSG levels induced by PDTC pro-oxidant activity, by restoring the intracellular redox equilibrium. Recent work, however, has shown that the effects of NAC on T cell apoptosis are not mediated by increased cellular glutathione (Jones *et al.*, 1995). NAC, in its capacity of a thiol, may also counter the pro-oxidant activity of PDTC more directly. That this may indeed be the case is strongly supported by recent observations made in our

laboratory showing that both L-cysteine and D-cysteine (which can not be incorporated into GSH) on their own are capable of blocking growth inhibition and apoptosis induced by PDTC in Tpi and Jurkat T cells (VH unpublished observation). Our observations also point towards a number of differences in the regulation of the $\text{I}\kappa\text{B}\alpha$ and $\text{I}\kappa\text{B}\beta$ degradation pathways. As has also previously been observed for Jurkat T cells and U937 monocytic cells (Suzuki *et al.*, 1994; Sen and Packer, 1996), OA-induced degradation of $\text{I}\kappa\text{B}\alpha$ could not be inhibited by PDTC in Tpi T cells. OA-induced degradation of $\text{I}\kappa\text{B}\beta$, on the other hand, was blocked (Figure 2B), demonstrating a differential sensitivity of the $\text{I}\kappa\text{B}\alpha$ and $\text{I}\kappa\text{B}\beta$ degradation pathways to PDTC-mediated inhibition. We also observed that OA was capable of inducing $\text{I}\kappa\text{B}\beta$ degradation in the transformed Tpi T cells, but not in resting LN T cells. This is reminiscent of the observations by Menon *et al.* (1993), who showed that OA treatment failed to induce nuclear translocation of NF- κB in several primary cell lines, whereas rapid induction by OA could readily be detected in their SV40-transformed counterparts. Both sets of observations support the concept that the regulation of $\text{I}\kappa\text{B}$ degradation pathways can differ depending on the state of activation of the cell.

NAC Blocks Apoptosis Induced by PDTC

Tpi and Jurkat T cells proved highly sensitive to PDTC and disulfiram, undergoing rapid cell cycle arrest and apoptosis. The fact that these processes are averted by NAC points towards different modes of action.

In other systems, PDTC and NAC either induce or inhibit apoptosis, depending on the cell line and the stimulus (Bessho *et al.*, 1994; Talley *et al.*, 1995; Kaneto *et al.*, 1996; Ivanov *et al.*, 1997). In cancer cells, PDTC can induce apoptosis and it also potentiates the activity of chemotherapeutic agents at doses that are not toxic to a range of untransformed epithelial cells (Chinery *et al.*, 1997). For this reason, clinical trials for the treatment of colorectal cancer and other solid tumours have been proposed, in which PDTC could be used in combination with chemotherapeutic agents (Chinery *et al.*, 1997). Our observations suggest that the potentiating, beneficial effects of PDTC during such treatment may not be possible in the presence of a drug such as NAC.

Interestingly, the anti-apoptotic effect of NAC is not specific for apoptosis induced by PDTC. In recent work, we demonstrated that NAC also blocks apoptosis induced by the alkylating agent N- α -tosyl-L-phenylalanine chloromethyl ketone (Heussler *et al.*, 1999a), which inhibits the NF- κB activation pathway in Tpi T cells by blocking $\text{I}\kappa\text{B}$ phosphorylation (Heussler *et al.*, 1999b).

Effects of NAC and PDTC on Other Signalling Pathways

AP-1, another key transcription factor, is also subject to redox regulation and its activation is also affected by PDTC and NAC (Meyer *et al.*, 1993; Munoz *et al.*, 1996;

Pinkus *et al.*, 1996; Yokoo and Kitamura, 1996). AP-1 transactivation is influenced by the intracellular thioredoxin and GSH status. PDTC, rather than inhibiting, induces AP-1-regulated gene expression. Such opposite effects on NF- κB and AP-1 activation have also been documented for H_2O_2 , thioredoxin and a number of other compounds (Meyer *et al.*, 1993; Galter *et al.*, 1994; Schenk *et al.*, 1994). Interestingly, Pinkus *et al.* (1996) demonstrated that PDTC-induced AP-1 activation is inhibited by pre-exposure of cells to GSH or antioxidants, including NAC. This is in line with our observations for the MAP kinase family members JNK and p38, and for the transcription factor ATF-2 which can be regulated by JNK and p38 (Gupta *et al.*, 1995; Raingeaud *et al.*, 1995; van Dam *et al.*, 1995; Wilhelm *et al.*, 1995). Treatment with PDTC enhanced JNK activation and this was blocked by NAC. In cells subjected to osmotic shock, JNK and p38 activity were enhanced by both NAC and PDTC, but strongly reduced in the presence of both components. NAC and PDTC, applied separately, did not interfere with ATF-2 activation induced by osmotic shock. On the contrary, PDTC tended to enhance ATF-2 phosphorylation. When osmotic shock was applied in the presence of both compounds, however, a clear reduction in the levels of phosphorylated ATF-2 could be observed. Our findings thus show that NAC blocks the effects exerted by PDTC on different signalling pathways, be they inhibitory, as is the case for NF- κB , or stimulatory, as was found for AP-1 and ATF-2. It has been reported that NF- κB activation can be regulated by MEKK-1 (Hirano *et al.*, 1996; Meyer *et al.*, 1996; Lee *et al.*, 1997a), a kinase which also controls the activation of JNK and p38. The physiological significance of MEKK1 vs. NIK or other kinases activators of IKK, however, has not yet been fully elucidated (Karin and Delhase, 1998).

Nevertheless, despite the fact that the NF- κB and JNK/p38 pathways may at one stage be regulated by a common upstream element, pathways further downstream clearly appear to differ in their sensitivity to NAC and PDTC (Pinkus *et al.*, 1996).

NF- κB represents a prime target for therapeutic intervention since blocking NF- κB activation inhibits both the production of inflammatory cytokines (such as TNF- α , IL1- β and IL6) by immune and endothelial cells (Munoz *et al.*, 1996) and the deleterious cellular changes induced by cytokines (Barnes and Adcock, 1997). NF- κB and AP-1 binding sites are also present in the promoter region of a number of genes that are directly involved in the pathogenesis of AIDS and cancer. The LTR of HIV-1 contains κB binding sites which may play a crucial role in regulating the latency of AIDS (Griffin *et al.*, 1989) and one of the major consequences of NF- κB activation is HIV gene transcription. Antioxidants such as NAC, dithiocarbamates and a wide variety of chemically unrelated inhibitors have been shown to block NF- κB activation or transcription directed by the HIV-LTR (Roederer *et al.*, 1990; Staal *et al.*, 1993) suggesting a possible therapeutic value in the clinical treatment of HIV-infected patients (Lemarie *et al.*, 1986; Reisinger *et al.*, 1990; Sunderman, 1991; Dröge *et al.*,

1992; Roederer *et al.*, 1992). As mentioned before, however, NAC can also enhance acute HIV-1 replication in monocyte-derived macrophages (Nottet *et al.*, 1997) and we now show that NAC can also counteract the inhibitory effects of dithiol compounds on HIV-LTR-driven reporter gene expression in T cells. In conclusion, it is becoming more and more obvious that NF- κ B responses, and the extent to which they can be inhibited, differ significantly according to cell type, stimulus or state of activation of the cell. At this stage, our observations are limited to NF- κ B activation in primary LN T cells and transformed T cells involving only a limited number of stimuli. The consistency with which NAC inhibited the effects of dithiocarbamates, however, suggests that it may be worthwhile testing whether these findings also apply to other systems. From a clinical point of view, our findings also provide a compelling argument for the judicious choice of drugs in the design of NF- κ B-based therapies.

Materials and Methods

Cell Lines and T Cell Cultures

The *T. parva*-infected (Tpi) T cell line TpM(803) is CD4⁺, CD8⁻, α/β TCR⁺ and its characteristics and maintenance have been described previously (Dobbelaere *et al.*, 1988). LN T cells were prepared as described previously (Mastro and Pepin, 1980).

Drug Treatment, Processing and Analysis

Chemical inhibitors or activators were added to the final concentrations as follows: cycloheximide (CHX; 177 nM; Sigma), pyrrolidine dithiocarbamate (PDTC; 100 μ M for short-term treatment in EMSA and 10 μ M for long-term treatment in transfection assays; Sigma), *N*-acetylcysteine (NAC; 30 mM or as indicated; Sigma), tetradecanoyl phorbol acetate (TPA; 3.3 nM; Sigma); ionomycin (2 μ M; Sigma); okadaic acid (OA; 2 or 4 μ M as indicated; LC laboratories); C2-ceramide (100 μ M or as indicated; Calbiochem, Lucerne, Switzerland); disulfiram (100 μ M for EMSA, 1 μ M for transfection assays or as indicated for proliferation assays; Sigma). Treatment of Tpi or LN T cells with NAC and/or PDTC was for 5 h; CHX treatment of LN T cells was for 5 h; for treatment of Tpi T cells, CHX was added for the last 4 h of NAC/PDTC treatment. Tpi T cells were induced with OA (2 μ M) for the last 90 min of NAC/PDTC treatment; LN T cells were induced with OA (4 μ M) for the last 40 min; disulfiram was added for the last 90 min. Ceramide stimulation was carried out for the last hour of NAC/PDTC treatment.

Proliferation assays were carried out as follows: 1×10^5 cells/ml were seeded in 96 flat-bottom well plates and cultured for 24 h in the presence or absence of NAC, PDTC or disulfiram, as indicated. Cells were pulsed with [methyl-³H]thymidine (0.2 μ Ci/well; 1 mCi = 37 MBq) for the last five hours and incorporation measured by liquid scintillation counting. Apoptosis was monitored using the Annexin-V-Fluos kit from Boehringer.

Cell Lysates and Immunoblot Analysis

Cells were collected and washed twice in ice-cold PBS, pH 7.4, and then processed for immunoblot analysis as described before (Palmer *et al.*, 1997). Rabbit anti-peptide polyclonal Ab with the following specificities were used: I κ B α (C-terminal 21 amino acids; sc-371, Santa Cruz, Basel, Switzerland); I κ B α (C-terminal

20 amino acids; sc-945); p105/p50 (C-terminal 20 amino acids; sc-114); RelA/p65 (C-terminal 20 amino acids; sc-372); ATF-2 (sc-187), phospho-c-Jun (SER63) II antibody (9261s, New England Biolabs, Allschwill, Switzerland) and phospho-specific p38 MAPK antibody (9211s, New England Biolabs). Bound primary Ab detected using horseradish peroxidase-conjugated goat anti-rabbit IgG and enhanced chemiluminescence (ECL, Amersham Corp).

Electrophoretic Mobility Shift Assay (EMSA), Cell Transfection and CAT Assays

EMSA was carried out using oligonucleotides (5'-AGTTGA-GGGGACTTTCCCAGGC-3' and its complement) in which the consensus NF- κ B binding site (underlined) is incorporated (Ivanov *et al.*, 1989). Cells were treated with NAC or PDTC for 5 h. OA (4 μ M) stimulation was carried out for 40 min, in the case of PDTC- or NAC-treated cells, during the last 40 min of treatment. Plasmids used for transfection have been described before (Ivanov *et al.*, 1989). Plasmid -121/+232 HIV-CAT and control -76/+232 HIV-CAT were derived by modification of pBLCAT5 containing the bacterial chloramphenicol acetyltransferase (CAT) gene. In plasmid -121/+232 HIV-CAT (abbreviated HIV-CAT), the CAT gene is under control of an HIV-1 enhancer with two κ B elements and the promoter region of the enhancer. Plasmid -76/+232 HIV-CAT (Δ HIV-CAT) is identical except for deletion of the region containing the κ B elements. Cells were transfected by electroporation using standard protocols (2×10^7 cells/ml, 600 μ l aliquots, 20 μ g of plasmid DNA, 430 V, 250 Ω). After electroporation, cells were placed on ice for 5 min and then diluted in Leibovitz L15 medium and incubated at 37 °C for 24 h. Treated or untreated cells were incubated for an additional 24 h, collected and washed with PBS, and then lysed by freeze-thawing in a CAT assay buffer (100 mM Tris, pH 7.4, 1 mM EDTA, 1 mM DTT). After centrifugation, the supernatant was collected, incubated for 6 min at 65 °C, then re-centrifuged. A 15 μ l aliquot was assayed for CAT activity using 9.25×10^3 Bq (250 nCi) of [¹⁴C]-chloramphenicol in CAT assay buffer supplemented with 100 μ g bovine serum albumin and 2 mg/ml *n*-butyryl coenzyme A. Butyrylated chloramphenicol was extracted using xylene and radioactivity determined by liquid scintillation counting.

Acknowledgements

This work was supported by the Swiss National Science Foundation (No. 31.43328.95 to D.D.) and Novartis (Ciba-Geigy Jubiläums-Stiftung), the Swiss Cancer League (KFS 625-2-1998), the US Department of Agriculture and Washington State University. Paula Fernandez is the recipient of a stipend (Swiss National Science Foundation No. 31.44407.95) for the MD-PhD-Programme.

References

- Albrecht, H., Tschopp, J., and Jongeneel, C.V. (1994). Bcl-2 protects from oxidative damage and apoptotic cell death without interfering with activation of NF- κ B by TNF. *FEBS Lett.* 357, 45–48.
- Anderson, M.T., Staal, F.J., Gitler, C., Herzenberg, L.A., and Herzenberg, L.A. (1994). Separation of oxidant-initiated and redox-regulated steps in the NF- κ B signal transduction pathway. *Proc. Natl. Acad. Sci. USA* 91, 11527–11531.

- Baeuerle, P.A., and Henkel, T. (1994). Function and activation of NF- κ B in the immune system. *Annu. Rev. Immunol.* **12**, 141–179.
- Baeuerle, P.A., and Baltimore, D. (1996). NF- κ B: ten years after. *Cell* **87**, 13–20.
- Baeuerle, P.A., Rupec, R.A., and Pahl, H.L. (1996). Reactive oxygen intermediates as second messengers of a general pathogen response. *Pathol. Biol. (Paris)* **44**, 29–35.
- Baldwin, A.S.J. (1996). The NF- κ B and I κ B proteins: new discoveries and insights. *Annu. Rev. Immunol.* **14**, 649–683.
- Ballou, L.R., Laulerkind, S.J., Rosloniec, E.F., and Raghov, R. (1996). Ceramide signalling and the immune response. *Biochim. Biophys. Acta* **1307**, 273–287.
- Barnes, P.J., and Adcock, I.M. (1997). NF- κ B: a pivotal role in asthma and a new target for therapy. *Trends Pharmacol. Sci.* **18**, 46–50.
- Bessho, R., Matsubara, K., Kubota, M., Kuwakado, K., Hirota, H., Wakazono, Y., Lin, Y.W., Okuda, A., Kawai, M., Nishikomori, R., and al, e. (1994). Pyrrolidine dithiocarbamate, a potent inhibitor of nuclear factor- κ B (NF- κ B) activation, prevents apoptosis in human promyelocytic leukemia HL-60 cells and thymocytes. *Biochem. Pharmacol.* **48**, 1883–1889.
- Blackwell, T.S., Blackwell, T.R., Holden, E.P., Christman, B.W., and Christman, J.W. (1996). *In vivo* antioxidant treatment suppresses nuclear factor- κ B activation and neutrophilic lung inflammation. *J. Immunol.* **157**, 1630–1637.
- Bowie, A., Moynagh, P.N., and O'Neill, L.A. (1996). Mechanism of NF- κ B activation by interleukin-1 and tumour necrosis factor in endothelial cells. *Biochem. Soc. Trans.* **24**, 2.
- Brennan, P., and O'Neill, L.A. (1995). Effects of oxidants and antioxidants on nuclear factor κ B activation in three different cell lines: evidence against a universal hypothesis involving oxygen radicals. *Biochim. Biophys. Acta* **1260**, 167–175.
- Brennan, P., and O'Neill, L.A. (1996a). 2-mercaptoethanol restores the ability of nuclear factor κ B (NF κ B) to bind DNA in nuclear extracts from interleukin 1-treated cells incubated with pyrrolidine dithiocarbamate (PDTC). Evidence for oxidation of glutathione in the mechanism of inhibition of NF- κ B by PDTC. *Biochem. J.* **320**, 975–981.
- Brennan, P., and O'Neill, L.A. (1996b). Inhibition of NF- κ B activity by oxidative processes in intact cells mechanism of action of pyrrolidine dithiocarbamate and diamide. *Biochem. Soc. Trans.* **24**, 3.
- Chinery, R., Brockman, J.A., Peeler, M.O., Shyr, Y., Beauchamp, R.D., and Coffey, R.J. (1997). Antioxidants enhance the cytotoxicity of chemotherapeutic agents in colorectal cancer: a p53-independent induction of p21WAF1/CIP1 via C/EBP β . *Nature Med.* **3**, 1233–1241.
- DiDonato, J., Mercurio, F., Rosette, C., Wu Li, J., Suyang, H., Ghosh, S., and Karin, M. (1996). Mapping of the inducible I κ B phosphorylation sites that signal its ubiquitination and degradation. *Mol. Cell Biol.* **16**, 1295–1304.
- DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E., and Karin, M. (1997). A cytokine-responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature* **388**, 548–554.
- Dobbelaere, D.A., Coquerelle, T.M., Roditi, I.J., Eichhorn, M., and Williams, R.O. (1988). *Theileria parva* infection induces autocrine growth of bovine lymphocytes. *Proc. Natl. Acad. Sci. USA* **85**, 4730–4734.
- Dröge, W., Eck, H.P., Gmunder, H., and Mihm, S. (1991). Modulation of lymphocyte functions and immune responses by cysteine and cysteine derivatives. *Am. J. Med.* **91**, 140–144.
- Dröge, W., Eck, H.P., and Mihm, S. (1992). HIV-induced cysteine deficiency and T-cell dysfunction – a rationale for treatment with N-acetylcysteine. *Immunol. Today* **13**, 211–214.
- Dröge, W., Schulze-Osthoff, K., Mihm, S., Galter, D., Schenk, H., Eck, H.P., Roth, S., and Gmunder, H. (1994). Functions of glutathione and glutathione disulfide in immunology and immunopathology. *FASEB J.* **8**, 1131–1138.
- Eichhorn, M., Prospero, T.D., Heussler, V.T., and Dobbelaere, D.A. (1993). Antibodies against major histocompatibility complex class II antigens directly inhibit the growth of T cells infected with *Theileria parva* without affecting their state of activation. *J. Exp. Med.* **178**, 769–776.
- Flohé, L., Brigelius-Flohé, R., Saliou, C., Traber, M.G., and Packer, L. (1997). Redox regulation of NF- κ B activation. *Free Radic. Biol. Med.* **22**, 1115–1126.
- Galley, Y., Hagens, G., Glaser, I., Davis, W.C., Eichhorn, M., and Dobbelaere, D.A.E. (1997). Jun NH2-terminal kinase is constitutively activated in T cells transformed by the intracellular parasite *Theileria parva*. *Proc. Natl. Acad. Sci. USA* **94**, 5119–5124.
- Galter, D., Mihm, S., and Dröge, W. (1994). Distinct effects of glutathione disulphide on the nuclear transcription factor κ B and the activator protein-1. *Eur. J. Biochem.* **221**, 639–648.
- Gazzinelli, R.T., Sher, A., Cheever, A., Gerstberger, S., Martin, M.A., and Dickie, P. (1996). Infection of human immunodeficiency virus 1 transgenic mice with *Toxoplasma gondii* stimulates proviral transcription in macrophages *in vivo*. *J. Exp. Med.* **183**, 1645–1655.
- Ginn Pease, M.E., and Whisler, R.L. (1996). Optimal NF κ B mediated transcriptional responses in Jurkat T cells exposed to oxidative stress are dependent on intracellular glutathione and costimulatory signals. *Biochem. Biophys. Res. Commun.* **226**, 695–702.
- Ginn Pease, M.E., and Whisler, R.L. (1998). Redox signals and NF- κ B activation in T cells. *Free Radic. Biol. Med.* **25**, 346–361.
- Giri, D.K., and Aggarwal, B.B. (1998). Constitutive activation of NF- κ B causes resistance to apoptosis in human cutaneous T cell lymphoma HuT-78 cells. *J. Biol. Chem.* **273**, 14008–14014.
- Gomez del Arco, P., Martinez Martinez, S., Calvo, V., Armesilla, A.L., and Redondo, J.M. (1996). JNK (c-Jun NH2-terminal kinase) is a target for antioxidants in T lymphocytes. *J. Biol. Chem.* **271**, 26335–26340.
- Griffin, G.E., Leung, K., Folks, T.M., Kunkel, S., and Nabel, G.J. (1989). Activation of HIV gene expression during monocyte differentiation by induction of NF- κ B. *Nature* **339**, 70–73.
- Gupta, S., Campbell, D., Derijard, B., and Davis, R.J. (1995). Transcription factor ATF2 regulation by the JNK signal transduction pathway. *Science* **267**, 389–393.
- Hellin, A.C., Calmant, P., Gielen, J., Bours, V., and Merville, M.P. (1998). Nuclear factor-kappaB-dependent regulation of p53 gene expression induced by daunomycin genotoxic drug. *Oncogene* **16**, 1187–1195.
- Heussler, V.T., Fernandez, P.C., Machado, J., Jr., Botteron, C., and Dobbelaere, D.A.E. (1999a). N-acetylcysteine blocks apoptosis induced by N- α -tosyl-L-phenylalanine chloromethyl ketone in transformed T cells. *Cell Death Diff.* **6**, 342–350.
- Heussler, V.T., Machado, J., Fernandez, P.C., Botteron, C., C.-G., C., Pearce, M.J., and Dobbelaere, D.A.E. (1999b). The intracellular parasite *Theileria parva* protects infected T cells from apoptosis. *Proc. Natl. Acad. Sci. USA* **96**, 7312–7317.
- Hirano, M., Osada, S., Aoki, T., Hirai, S., Hosaka, M., Inoue, J., and Ohno, S. (1996). MEK kinase is involved in tumor necrosis factor α -induced NF- κ B activation and degradation of I κ B- α . *J. Biol. Chem.* **271**, 13234–13238.
- Israel, N., Gougerot Pocidalo, M.A., Aillet, F., and Virelizier, J.L. (1992). Redox status of cells influences constitutive or induced NF- κ B translocation and HIV long terminal repeat activity

- in human T and monocytic cell lines. *J. Immunol.* **149**, 3386–3393.
- Ivanov, V., Stein, B., Baumann, I., Dobbelaere, D.A., Herrlich, P., and Williams, R.O. (1989). Infection with the intracellular protozoan parasite *Theileria parva* induces constitutively high levels of NF- κ B in bovine T lymphocytes. *Mol. Cell Biol.* **9**, 4677–4686.
- Ivanov, V.N., Lee, R.K., Podack, E.R., and Malek, T.R. (1997). Regulation of Fas-dependent activation-induced T cell apoptosis by cAMP signaling: a potential role for transcription factor NF- κ B. *Oncogene* **14**, 2455–2464.
- Jones, D.P., Maellaro, E., Jiang, S., Slater, A.F., and Orrenius, S. (1995). Effects of N-acetyl-L-cysteine on T cell apoptosis are not mediated by increased cellular glutathione. *Immunol. Lett.* **45**, 205–209.
- Kaneto, H., Fujii, J., Myint, T., Miyazawa, N., Islam, K.N., Kawasaki, Y., Suzuki, K., Nakamura, M., Tatsumi, H., Yamasaki, Y., and Taniguchi, N. (1996). Reducing sugars trigger oxidative modification and apoptosis in pancreatic beta-cells by provoking oxidative stress through the glycation reaction. *Biochem. J.* **320**, 855–863.
- Karin, M., and Delhase, M. (1998). JNK or IKK, AP-1 or NF- κ B, which are the targets for MEK kinase 1 action? *Proc. Natl. Acad. Sci. USA* **95**, 9067–9069.
- Kikumori, T., Kambe, F., Nagaya, T., Imai, T., Funahashi, H., and Seo, H. (1998). Activation of transcriptionally active nuclear factor- κ B by tumor necrosis factor- α and its inhibition by antioxidants in rat thyroid FRTL-5 cells. *Endocrinology* **139**, 1715–1722.
- Krappmann, D., Wulczyn, F.G., and Scheidereit, C. (1996). Different mechanisms control signal-induced degradation and basal turnover of the NF- κ B inhibitor I κ B alpha *in vivo*. *EMBO J.* **15**, 6716–6726.
- Lee, F.S., Hagler, J., Chen, Z.J., and Maniatis, T. (1997a). Activation of the I κ B- α kinase complex by MEKK1, a kinase of the JNK pathway. *Cell* **88**, 213–222.
- Lee, R., Beauparlant, P., Elford, H., Ponka, P., and Hiscott, J. (1997b). Selective inhibition of I κ B- α phosphorylation and HIV-1 LTR-directed gene expression by novel antioxidant compounds. *Virology* **234**, 277–290.
- Legrand Poels, S., Zecchinon, L., Piret, B., Schoonbroodt, S., and Piette, J. (1997). Involvement of different transduction pathways in NF- κ B activation by several inducers. *Free Radic. Res.* **27**, 301–309.
- Lemarie, E., Musset, M., Charbonnier, C., Renoux, M., and Renoux, G. (1986). Clinical characterization of imuthiol. *Methods Find. Exp. Clin. Pharmacol.* **8**, 51–54.
- Li, N., and Karin, M. (1999). Is NF- κ B the sensor of oxidative stress? *FASEB J.* **13**, 1137–1143.
- Mastro, A., and Pepin, K. (1980). Suppression of lectin-stimulated DNA synthesis in bovine lymphocytes by the tumor promoter 12-O-tetradecanoylphorbol-13-acetate. *Cancer Res.* **40**, 3307–3312.
- Menon, S.D., Qin, S., Guy, G.R., and Tan, Y.H. (1993). Differential induction of nuclear NF- κ B by protein phosphatase inhibitors in primary and transformed human cells. Requirement for both oxidation and phosphorylation in nuclear translocation. *J. Biol. Chem.* **268**, 26805–26812.
- Menon, S.D., Guy, G.R., and Tan, Y.H. (1995). Involvement of a putative protein-tyrosine phosphatase and I κ B- α serine phosphorylation in nuclear factor- κ B activation by tumor necrosis factor. *J. Biol. Chem.* **270**, 18881–18887.
- Meyer, M., Schreck, R., and Baeuerle, P.A. (1993). H₂O₂ and antioxidants have opposite effects on activation of NF- κ B and AP-1 in intact cells: AP-1 as secondary antioxidant-responsive factor. *EMBO J.* **12**, 2005–2015.
- Meyer, M., Pahl, H.L., and Baeuerle, P.A. (1994). Regulation of the transcription factors NF- κ B and AP-1 by redox changes. *Chem. Biol. Interact.* **97**, 91–100.
- Meyer, C.F., Wang, X., Chang, C., Templeton, D., and Tan, T.H. (1996). Interaction between c-Rel and the mitogen-activated protein kinase kinase kinase 1 signaling cascade in mediating κ B enhancer activation. *J. Biol. Chem.* **271**, 8971–8976.
- Mihm, S., Galter, D., and Dröge, W. (1995). Modulation of transcription factor NF- κ B activity by intracellular glutathione levels and by variations of the extracellular cysteine supply. *FASEB J.* **9**, 246–252.
- Miyamoto, S., Maki, M., Schmitt, M.J., Hatanaka, M., and Verma, I.M. (1994). Tumor necrosis factor α -induced phosphorylation of I κ B- α is a signal for its degradation but not dissociation from NF- κ B. *Proc. Natl. Acad. Sci. USA* **91**, 12740–12744.
- Munoz, C., Pascual Salcedo, D., Castellanos, M.C., Alfranca, A., Aragonés, J., Vara, A., Redondo, M.J., and de Landazuri, M.O. (1996). Pyrrolidine dithiocarbamate inhibits the production of interleukin-6, interleukin-8, and granulocyte-macrophage colony-stimulating factor by human endothelial cells in response to inflammatory mediators: modulation of NF- κ B and AP-1 transcription factors activity. *Blood* **88**, 3482–3490.
- Nagata, S. (1997). Apoptosis by death factor. *Cell* **88**, 355–365.
- Nobel, C.I., Kimland, M., Lind, B., Orrenius, S., and Slater, A.F. (1995). Dithiocarbamates induce apoptosis in thymocytes by raising the intracellular level of redox-active copper. *J. Biol. Chem.* **270**, 26202–26208.
- Nottet, H.S., Moelans, I.I., de Vos, N.M., de Graaf, L., Visser, M.R., and Verhoef, J. (1997). N-acetyl-L-cysteine-induced up-regulation of HIV-1 gene expression in monocyte-derived macrophages correlates with increased NF- κ B DNA binding activity. *J. Leukoc. Biol.* **61**, 33–39.
- Palmer, G.H., Machado, J.J., Fernandez, P., Heussler, V., Perinat, T., and Dobbelaere, D.A. (1997). Parasite-mediated nuclear factor- κ B regulation in lymphoproliferation caused by *Theileria parva* infection. *Proc. Natl. Acad. Sci. USA* **94**, 12527–12532.
- Palombella, V.J., Rando, O.J., Goldberg, A.L., and Maniatis, T. (1994). The ubiquitin-proteasome pathway is required for processing the NF- κ B1 precursor protein and the activation of NF- κ B. *Cell* **78**, 773–785.
- Pinkus, R., Weiner, L.M., and Daniel, V. (1996). Role of oxidants and antioxidants in the induction of AP-1, NF- κ B, and glutathione S-transferase gene expression. *J. Biol. Chem.* **271**, 13422–13429.
- Poli, G., and Fauci, A.S. (1992). The effect of cytokines and pharmacologic agents on chronic HIV infection. *AIDS Res. Hum. Retroviruses* **8**, 191–197.
- Raingaud, J., Gupta, S., Rogers, J.S., Dickens, M., Han, J., Ulevitch, R.J., and Davis, R.J. (1995). Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J. Biol. Chem.* **270**, 7420–7426.
- Reisinger, E.C., Kern, P., Ernst, M., Bock, P., Flad, H.D., and Dietrich, M. (1990). Inhibition of HIV progression by dithiocarb. German DTC Study Group. *Lancet* **335**, 679–682.
- Rieckmann, P., Thevenin, C., and Kehrl, J.H. (1992). Okadaic acid is a potent inducer of AP-1, NF- κ B, and tumor necrosis factor- α in human B lymphocytes. *Biochem. Biophys. Res. Commun.* **187**, 51–57.
- Roederer, M., Ela, S.W., Staal, F.J., Herzenberg, L.A., and Herzenberg, L.A. (1992). N-acetylcysteine: a new approach to anti-HIV therapy. *AIDS Res. Hum. Retroviruses* **8**, 209–217.
- Roederer, M., Staal, F.J., Raju, P.A., Ela, S.W., Herzenberg, L.A., and Herzenberg, L.A. (1990). Cytokine-stimulated human immunodeficiency virus replication is inhibited by N-acetyl-L-cysteine. *Proc. Natl. Acad. Sci. USA* **87**, 4884–4888.

- Sakurada, S., Kato, T., and Okamoto, T. (1996). Induction of cytokines and ICAM-1 by proinflammatory cytokines in primary rheumatoid synovial fibroblasts and inhibition by N-acetyl-L-cysteine and aspirin. *Int. Immunol.* *8*, 1483–1493.
- Schenk, H., Klein, M., Erdbrugger, W., Dröge, W., and Schulze-Osthoff, K. (1994). Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF- κ B and AP-1. *Proc. Natl. Acad. Sci. USA* *91*, 1672–1676.
- Schofield, L., Novakovic, S., Gerold, P., Schwarz, R.T., McConville, M.J., and Tachado, S.D. (1996). Glycosylphosphatidylinositol toxin of Plasmodium up-regulates intercellular adhesion molecule-1, vascular cell adhesion molecule-1, and E-selectin expression in vascular endothelial cells and increases leukocyte and parasite cytoadherence via tyrosine kinase-dependent signal transduction. *J. Immunol.* *156*, 1886–1896.
- Schreck, R., Rieber, P., and Baeuerle, P.A. (1991). Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF- κ B transcription factor and HIV-1. *EMBO J.* *10*, 2247–2258.
- Schreck, R., Meier, B., Mannel, D.N., Dröge, W., and Baeuerle, P.A. (1992). Dithiocarbamates as potent inhibitors of nuclear factor- κ B activation in intact cells. *J. Exp. Med.* *175*, 1181–1194.
- Schreck, R., and Baeuerle, P.A. (1994). Assessing oxygen radicals as mediators in activation of inducible eukaryotic transcription factor NF- κ B. *Methods Enzymol.* *234*, 151–163.
- Schulze-Osthoff, K., Bauer, M.K., Vogt, M., and Wesselborg, S. (1997). Oxidative stress and signal transduction. *Int. J. Vitam. Nutr. Res.* *67*, 336–342.
- Schulze-Osthoff, K., Beyaert, R., Vandevorde, V., Haegeman, G., and Fiers, W. (1993). Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF. *EMBO J.* *12*, 3095–3104.
- Sen, C.K., and Packer, L. (1996). Antioxidant and redox regulation of gene transcription. *FASEB J.* *10*, 709–720.
- Siebenlist, U., Franzoso, G., and Brown, K. (1994). Structure, regulation and function of NF- κ B. *Annu. Rev. Cell Biol.* *10*, 405–455.
- Staal, F.J., Roederer, M., Raju, P.A., Anderson, M.T., Ela, S.W., Herzenberg, L.A., and Herzenberg, L.A. (1993). Antioxidants inhibit stimulation of HIV transcription. *AIDS Res. Hum. Retroviruses* *9*, 299–306.
- Stancovski, I., and Baltimore, D. (1997). NF- κ B activation: the I κ B kinase revealed? *Cell* *91*, 299–302.
- Sunderman, F.W., Sr. (1991). Therapeutic properties of sodium diethyldithiocarbamate: its role as an inhibitor in the progression of AIDS. *Ann. Clin. Lab. Sci.* *21*, 70–81.
- Suzuki, Y.J., Mizuno, M., and Packer, L. (1994). Signal transduction for nuclear factor- κ B activation. Proposed location of antioxidant-inhibitable step. *J. Immunol.* *153*, 5008–5015.
- Suzuki, Y.J., Mizuno, M., and Packer, L. (1995). Transient overexpression of catalase does not inhibit TNF- or PMA- induced NF- κ B activation. *Biochem. Biophys. Res. Commun.* *210*, 537–541.
- Talley, A.K., Dewhurst, S., Perry, S.W., Dollard, S.C., Gummuluru, S., Fine, S.M., New, D., Epstein, L.G., Gendelman, H.E., and Gelbard, H.A. (1995). Tumor necrosis factor α -induced apoptosis in human neuronal cells: protection by the antioxidant N-acetylcysteine and the genes bcl-2 and crmA. *Mol. Cell Biol.* *15*, 2359–2366.
- Traenckner, E.B., Wilk, S., and Baeuerle, P.A. (1994). A proteasome inhibitor prevents activation of NF- κ B and stabilizes a newly phosphorylated form of I κ B- α that is still bound to NF- κ B. *EMBO J.* *13*, 5433–5441.
- van Dam, H., Wilhelm, D., Herr, I., Steffen, A., Herrlich, P., and Angel, P. (1995). ATF-2 is preferentially activated by stress-activated protein kinases to mediate c-jun induction in response to genotoxic agents. *EMBO J.* *14*, 1798–1811.
- Verhoven, B., Schlegel, R.A., and Williamson, P. (1995). Mechanisms of phosphatidylserine exposure, a phagocyte recognition signal, on apoptotic T lymphocytes. *J. Exp. Med.* *182*, 1597–1601.
- Verma, I.M., Stevenson, J.K., Schwarz, E.M., Van Antwerp, D., and Miyamoto, S. (1995). Rel/NF- κ B/I κ B family: intimate tales of association and dissociation. *Genes Dev.* *9*, 2723–2735.
- Whitmarsh, A.J., and Davis, R.J. (1996). Transcription factor AP-1 regulation by mitogen-activated protein kinase signal transduction pathways. *J. Mol. Med.* *74*, 589–607.
- Wilhelm, D., van Dam, H., Herr, I., Baumann, B., Herrlich, P., and Angel, P. (1995). Both ATF-2 and c-Jun are phosphorylated by stress-activated protein kinases in response to UV irradiation. *Immunobiology* *193*, 143–148.
- Yokoo, T., and Kitamura, M. (1996). Antioxidant PDTC induces stromelysin expression in mesangial cells via a tyrosine kinase-AP-1 pathway. *Am. J. Physiol.* *270*, 806–811.

Received July 7, 1999; accepted October 27, 1999