Genetic variability and limited clonality of *Mycoplasma hyorhinis* in pig herds

Bettina Trüeb^{a,1}, Elena Catelli^{b,1}, Adrian Luehrs^c, Heiko Nathues^d, Peter Kuhnert^{a,*}

^a*Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Laenggassstr. 122, 3001 Bern, Switzerland*

^b*Department of Veterinary Science, University of Parma, Via del Taglio 8, 43126 Parma, Italy*

^c*Field Station for Epidemiology, University of Veterinary Medicine Hannover, Foundation, Bueschelerstr. 9, 49456 Bakum, Germany*

^d*Clinic for Swine, Department of Clinical Veterinary Medicine, Vetsuisse Faculty, University of Bern, Bremgartenstr. 109a, 3001 Bern, Switzerland*

¹equal contribution

*Corresponding author: Peter Kuhnert, Institute of Veterinary Bacteriology, University of Bern, Laenggassstr. 122, 3001 Bern, Switzerland. peter.kuhnert@vetsuisse.unibe.ch

Tel +41 31 6312485

Fax +41 31 6312634

21 **Abstract**

22 *Mycoplasma hyorhinis* is a common inhabitant of the upper respiratory tract and tonsils of pigs. Its
23 role as a possible pathogen remains controversial. In order to gain more insight into the
24 epidemiology and population structure of *Mycoplasma hyorhinis* we genetically characterized 60
25 isolates by multi locus sequence typing (MLST). The *M. hyorhinis* strains originated from Swiss
26 and German pig herds with knowledge on the clinical background. The MLST scheme of
27 Tocqueville *et al.* (J.Clin.Microbiol. 2014) was optimized, primers for the six MLST gene
28 fragments were newly designed to allow amplification and sequencing with a single protocol. A
29 total of 27 ST were observed with the 60 strains, 26 of those were previously unknown types.
30 Generally identical genotypes were observed within a farm but they differed between farms. The
31 identical genotype was also observed in three different Swiss farms. On the other hand different
32 genotypes within a farm were found with three German farms. The Swiss isolates formed a distinct
33 cluster but otherwise there was no geographical nor a clinical association with specific clusters
34 observed. Data shows a high variability of *M. hyorhinis* comparable to what is observed for
35 *Mycoplasma hyopneumoniae*. Similar to this pathogen the population structure of *M. hyorhinis* also
36 shows some limited clonality with predominant genotypes within an animal and a single farm but
37 different ones between farms. The comparable population structure of *M. hyopneumoniae* and *M.*
38 *hyorhinis* could indicate a similar evolution of the two species in the common pig host.

39

40 Keywords: MLST, genotyping, pig disease, enzootic pneumonia, coughing

41

42 **1. Introduction**

43 *Mycoplasma hyorhinis*, a common inhabitant of the upper respiratory tract and tonsils of pigs, was
44 first described in the fifties by Switzer (1955). It is generally regarded as a commensal, however, a
45 role of *M. hyorhinis* as a potential pathogen has also been discussed. Infections with this organism
46 are subclinical, even though arthritis, polyserositis, conjunctivitis and otitis as well as abortions are
47 clinical disorders that may be observed in an infected pig (Friis and Feenstra, 1994; Kobisch and
48 Friis, 1996; Shin et al., 2003). *Mycoplasma hyorhinis* has also been found associated with
49 pneumonia, however, rather as a secondary pathogen (Falk et al., 1991; Kawashima et al., 1996;
50 Kobayashi et al., 1996; Lin et al., 2006). Nevertheless, an increase in the detection of *M. hyorhinis*
51 in recent years has been observed and the relevance of this pathogen as an emerging issue causing
52 significant economic losses is suspected (Dos Santos et al., 2015).

53 Genotyping allows epidemiological investigation and population studies. This can help to clarify
54 the role of a possible pathogen, identify specifically involved strains and their transmission.
55 Multilocus sequence typing (MLST) is a highly selective and specific method which has been
56 utilized for the characterization of many bacterial species including *Mycoplasma hyopneumoniae*
57 and *M. hyorhinis* (Mayor et al., 2008; Tocqueville et al., 2014). Genotyping of *M. hyorhinis* using
58 MLST is based on sequences of internal fragments from six housekeeping genes, i.e. the
59 chromosomal replication initiation protein (*dnaA*), the RNA polymerase β subunit (*rpoB*), the DNA
60 gyrase subunit B (*gyrB*), the glutamyl-tRNA synthetase (*gltX*), the adenylate kinase (*adk*) and the
61 guanylate kinase (*gmk*).

62 In this study, we optimized the MLST scheme of six loci by defining completely new primer sets.
63 They were used to genetically characterize 60 *M. hyorhinis* isolates belonging to twenty-five Swiss
64 and German herds. Information about the herd health status, i.e. clinical respiratory disease, of the
65 fattening units from where strains came from were gathered as well. The MLST data of the
66 *M. hyorhinis* isolates was used to elucidate genetic variability of the species and was related to the
67 clinical presentation of the pig in order to define possible virulent strains.

68 **2. Materials and methods**

69 *2.1. Strains and template preparation*

70 A total of 60 *M. hyorhinis* isolates with 47 from Germany and 13 from Switzerland previously
71 included in a transnational cross sectional study determining the prevalence of *M. hyorhinis* were
72 analyzed (Luehrs, 2015). For every source of strains, relevant pathological conditions in the
73 corresponding animals were available from databases. This includes presence of (enzootic)
74 pneumonia varying from mild, moderate to severe assessed during lung scoring after dissection, as
75 well as arthritis and/or polyserositis. Moreover, the simultaneous infection with *M. hyopneumoniae*
76 was examined in every sample using real-time PCR (Dubosson et al., 2004).

77 Strains of *M. hyorhinis* were kept at -80°C and cultured fresh in 1 ml liquid medium (Mycoplasma
78 Experience, Bletchingley, Great Britain) for 1-2 days at 37°C. Once medium color change from red
79 to yellow occurred the culture was centrifuged for 10 min. at 5000xg. The pellet was dissolved in
80 70µl ddH₂O and boiled for 15 min. at 95°C. This crude lysate was then kept at -20°C until being
81 used for PCR.

82

83 *2.2. Optimization of MLST*

84 The MLST scheme of Tocqueville et al. (2014) using the six housekeeping genes *dnaA*, *rpoB*, *gyrB*,
85 *gltX*, *adk* and *gmk* served as a basis to optimize the protocol. Primers with identical annealing
86 temperatures were defined, allowing amplification of all targets with a single PCR protocol instead
87 of 3 different ones as described by Tocqueville et al. (2014). For this purpose target genes were
88 extracted from available genome sequences of *M. hyorhinis* strains DBS 1050, HUB-1, GDL-1,
89 MCLD, SK76 as well as the type strain ATCC17981^T (=BTS7^T) and sequences were aligned.
90 Consensus sequence stretches as close as possible to the starting sequence used for typing were
91 defined and primers with similar annealing temperature selected (Table 1). Each PCR was
92 performed in a 30-µl total volume containing 12 pmol of each primer, 1 x FIREPol® Master Mix
93 Ready to Load (Solis BioDyne, Tartu, Estonia) and 1 µl lysate at the following cycling conditions:

94 3 min denaturation at 95°C, followed by 35 cycles at 95°C for 30 s, 55°C for 30 s and 72°C for 45
95 s. A final extension step for 7 min at 72°C was included. Subsequently, the amplification products
96 of the six genes from one strain were pooled and then purified using the “High pure PCR
97 purification Kit” (Roche Diagnostics, Rotkreuz, Switzerland). Finally the purified pooled PCR
98 products was used for sequencing with the BigDye® Terminator v3.1 Cycle Sequencing Kit
99 (Thermo Fisher, Reinach, Switzerland), using the same primers as for PCR (Table 1). For this
100 purpose 96-well plates (Thermo Fisher) were prepared with corresponding primers (5 pmol/well)
101 using a multichannel pipette. The plates allowed batch-wise analysis of 8 strains and could be kept
102 at -20°C until being used for sequencing (Korczak et al., 2009).

103

104 2.3. Data analysis

105 Sequences were edited using the Sequencher program v5.1 (GeneCodes, Ann Arbor, MI, USA) and
106 imported into Bionumerics v7.5 (AppliedMaths, Sint Marten, Belgium). Sequence types (ST) were
107 determined using the MLST plugin allowing direct query of the PubMLST database
108 (www.pubmlst.org). New alleles and allele combinations were submitted to the curator of the
109 database to assign allele and ST numbers. The MLST sequences of the 33 strains from the study of
110 Tocqueville et al. (2014) as well as the ones of the type strain BTS7^T (GenBank accession
111 ARTL000000000) were also imported to the database and included in the analyses. The resulting 94
112 entries were used for creating a minimal spanning tree based on allele numbers and a UPGMA tree
113 based on allele sequences in Bionumerics.

114 The START2 program (pubmlst.org) was used to calculate the ratio of synonymous and
115 nonsynonymous substitutions as well as the index of association (I_A). The index of discrimination
116 (D) was calculated using the formula of Hunter and Gaston (1988). LIAN
117 (guanine.evolbio.mpg.de/cgi-bin/lian/lian.cgi.pl) was used to calculate the genetic diversity (H).

118

119

120 3. Results

121 3.1. Genetic diversity of *M. hyorhinis*

122 All 60 isolates could be analyzed with the optimized MLST scheme. There were two new alleles
123 observed for *dnaA* (totaling 12) and *gyrB* (totaling 8), as well as three new alleles for the targets
124 *rpoB* (totaling 10), *gltX* (totaling 10), *adk* (totaling 7) and *gmk* (totaling 8). With the additional new
125 alleles the genetic diversity (H) of the six loci varies from 0.4155 (*gltX*) to 0.7596 (*gyrB*). The ratios
126 of nonsynonymous and synonymous substitutions varied from 0.1303 (*dnaA*) to 0.6785 (*gmk*). The
127 index of association (I_A) was 0.026, indication no linkage disequilibrium. A total of 27 ST were
128 defined, 26 of them were new resulting in a total of 55 different *M. hyorhinis* genotypes based on
129 MLST. Including all strains, i.e. also those from the same herd and same animals, the index of
130 discrimination (D) was 0.982. When only single representatives of such repetitive isolations were
131 included in the analysis the index of discrimination (D) was 0.995.

132 All the isolates from our study belong to the clonal complex 1 as defined by at least 4 common
133 alleles (Tocqueville et al., 2014). The clonal relationship of STs is graphically shown by the
134 minimal spanning tree in Fig. 1. Six of the seven previously observed singletons remained while
135 ST-28 was integrated into complex 1 (Tocqueville et al., 2014). There is only one common ST (ST-
136 13) between our study and the latter one as shown in Fig.2.

137

138 3.2. Limited clonality of *M. hyorhinis*

139 There are no clear and discrete geographic clusters present, even though some grouping can be
140 observed. The type strain BTS7^T originating from the USA was determined as ST-16 based on *in*
141 *silico* analysis of its genome sequence. This ST was also observed in France. The Chinese isolate
142 represents an out-group while most of the branches consist of isolates from more than one country.
143 Noteworthy, all Swiss isolates can be found on a single cluster containing only isolates from one
144 German farm where polyserositis was diagnosed in piglets. Another discernable cluster is formed
145 by only German isolates. This cluster contains isolates from farm DE_03 where two slightly

different strains based on a single base-pair change in *gmk* could be detected. Clearly different strains were observed in farm DE_16. In all other cases isolates from a single farm had the same genotype, while isolates from different farms were also of different genotype. Identical genotypes were also found when multiple isolates from single animals were analyzed (indicated by C2, C3). Identical genotypes were observed in three Swiss farms. In German farms, *M. hyorhinis* was isolated from animals that were in most cases (32 of 44) affected by *M. hyopneumoniae* as was diagnosed by real-time PCR indicated by ABC/REP in Fig. 2. In one case with a clear clinical status of enzootic pneumonia (strain numbers 10003) a follow up was done a few months later (strain numbers. 10025). While at both stages *M. hyorhinis* could be isolated from broncho-alveolar lavage fluid (BALF) and showed identical genotypes in different animals, the isolates from the two different time points differed in genotypes, indicating a switch of *M. hyorhinis* population over time.

There were no clusters or genotypes associated with certain clinical manifestations.

4. Discussion

This study analyzed the genetic variability of *M. hyorhinis* from pig farms in Germany and Switzerland. Genotypes were put in relation to available clinical data and were also compared to isolates contained in the PubMLST database. For this purpose the MLST scheme of Tocqueville et al. (2014) was optimized to allow a uniform PCR amplification and sequencing protocol. The new protocol, based on a completely new set of primers for the six target genes, proved to be robust. All 60 isolates could be genotyped. The genetic diversity of *M. hyorhinis* increased by 26 genotypes to a total of 55 different STs based on MLST. While in the previous study by Tocqueville et al. (2014) mainly isolates from France were genotyped, we analyzed strains from Germany and Switzerland. Except from one farm, where the previously identified ST-13 was found, all other were new STs. Generally the same ST was observed within a farm and also within a single animal. The same ST

172 could be observed even within three different Swiss farms. On the other side different ST were
173 observed in three German farms (DE_16, DE_03 and DE_05). From farm DE_05 one sample
174 resulted in an ambiguous base call at two positions what made a genotyping impossible but showed
175 that it contained a mix of at least two different strains (data not shown). This was an indirect
176 indication that indeed more than one strain can be present in a single animal, even though it might
177 not be the rule. Nevertheless, the observation that within a farm and individual animal the same
178 genotype is generally found is a fact that would allow for direct genotyping of clinical material.
179 This approach is successfully used with *M. hyopneumoniae* (Kuhnert and Overesch, 2014; Kuhnert
180 et al., 2011; Mayor et al., 2008) and could therefore also be applied for *M. hyorhinis*. This allows a
181 rapid straight forward genotyping eliminating the culture step.

182 The high variability but still limited clonality of *M. hyorhinis* is very similar to what can be
183 observed with *M. hyopneumoniae* (Kuhnert and Overesch, 2014; Mayor et al., 2008; Mayor et al.,
184 2007; Nathues et al., 2011). Given this observation and fact that *M. hyorhinis* is mainly a
185 commensal inhabiting the same niche as the more pathogenic *M. hyopneumoniae* could point to a
186 similar selection pressure if not co-evolution of the two species. Nevertheless, while whole genome
187 comparisons between these two species showed that rearrangements are common, clear differences
188 in gene composition indicate that a direct interaction and dependency of the two is rather unlikely.

189 Clinical information was gathered from all cases including diagnostic results on enzootic
190 pneumonia (EP) caused by *M. hyopneumoniae*. While *M. hyopneumoniae* was detected in more
191 than 70% of German cases, all Swiss cases were negative for this pathogen (Fig.2). This reflects the
192 different situation in these two countries. While in Switzerland only a few sporadic cases of EP are
193 observed per year due to a successful eradication program, EP is still endemic in Germany (Nathues
194 et al., 2014; Stark et al., 2007). Furthermore, clinical data is also available for strains contained in
195 the PubMLST database (Tocqueville et al., 2014). No correlation or association of specific
196 genotypes or clusters with clinical manifestations could be discerned. Similar observations were
197 made by Dos Santos et al. (2015) using a multi-locus variable-number tandem-repeat analysis

198 (MLVA) that is based on more variable genes than the house-keeping genes used for MLST. These
199 authors also could not find an association between isolates and clinical manifestations. The MLVA
200 approach was less discriminating than MLST with a diversity index of 0.814 compared to 0.982,
201 respectively. However, the MLVA is based on only 2 genes compared to the six genes used for
202 MLST and is therefore less work intensive and less expensive than the MLST. Nevertheless, the
203 MLST is highly reproducible with all six targets successfully analyzable in all strains whereas with
204 MLVA one of the targets could not be amplified in about 2% of strains and more than 40% of
205 clinical specimens (Dos Santos et al., 2015). Moreover, we could improve the MLST by designing
206 new primers that allow a single PCR amplification and sequencing protocol. The PCR purification
207 step using pooled samples as well as the pre-preparing of 96-well microtiter plates containing all
208 sequencing primers for 8 strains further reduce hands on time and costs. This makes MLST the
209 method of choice if comparing efforts and outcome.

210 In conclusion the MLST scheme for *M. hyorhinis* was optimized and the number of isolates
211 analyzed by this method was extended. Thereby the variability of *M. hyorhinis* in farms and related
212 clinical data of the host to genetic data of the potential pathogen was systematically investigated. A
213 limited clonality of *M. hyorhinis* was observed, i.e. a high variability of genotypes was
214 accompanied by identical genotypes within a farm and individual animal. No specifically virulent
215 genotypes could be discerned, indicating the commensal nature of *M. hyorhinis* in the pig lung.

216

217

218 **5. Acknowledgements**

219 This project was partly funded by the European Union's Horizon 2020 Programme under the Grant
220 Agreement n°633184 and Swiss Government.

221

222 **6. References**

223

- 224 Dos Santos, L.F., Clavijo, M.J., Sreevatsan, S., Rovira, A., Moreira, M.A., Pieters, M., 2015.
225 Genotyping of *Mycoplasma hyorhinis* using multiple-locus variable number tandem repeat
226 analysis. *J. Microbiol. Methods* 111, 87-92.
- 227 Dubosson, C.R., Conzelmann, C., Miserez, R., Boerlin, P., Frey, J., Zimmermann, W., Häni, H.,
228 Kuhnert, P., 2004. Development of two real-time PCR assays for the detection of
229 *Mycoplasma hyopneumoniae* in clinical samples. *Vet. Microbiol.* 102, 55-65.
- 230 Falk, K., Hoie, S., Lium, B.M., 1991. An abattoir survey of pneumonia and pleuritis in slaughter
231 weight swine from 9 selected herds. II. Enzootic pneumonia of pigs: microbiological
232 findings and their relationship to pathomorphology. *Acta Vet. Scand.* 32, 67-77.
- 233 Friis, N.F., Feenstra, A.A., 1994. *Mycoplasma hyorhinis* in the etiology of serositis among piglets.
234 *Acta Vet. Scand.* 35, 93-98.
- 235 Hunter, P.R., Gaston, M.A., 1988. Numerical index of the discriminatory ability of typing systems:
236 an application of Simpson's index of diversity. *J. Clin. Microbiol.* 26, 2465-2466.
- 237 Kawashima, K., Yamada, S., Kobayashi, H., Narita, M., 1996. Detection of porcine reproductive
238 and respiratory syndrome virus and *Mycoplasma hyorhinis* antigens in pulmonary lesions of
239 pigs suffering from respiratory distress. *J. Comp. Pathol.* 114, 315-323.
- 240 Kobayashi, H., Morozumi, T., Miyamoto, C., Shimizu, M., Yamada, S., Ohashi, S., Kubo, M.,
241 Kimura, K., Mitani, K., Ito, N., Yamamoto, K., 1996. *Mycoplasma hyorhinis* infection
242 levels in lungs of piglets with porcine reproductive and respiratory syndrome (PRRS). *J.*
243 *Vet. Med. Sci.* 58, 109-113.
- 244 Kobisch, M., Friis, N.F., 1996. Swine mycoplasmoses. *Rev. Sci. Tech.* 15, 1569-1605.
- 245 Korczak, B.M., Zurfluh, M., Emler, S., Kuhn-Oertli, J., Kuhnert, P., 2009. Multiplex strategy for
246 MLST, *fla*-typing and genetic determination of antimicrobial resistance of Swiss
247 *Campylobacter jejuni* and *Campylobacter coli* isolates. *J. Clin. Microbiol.* 47, 1996-2007.

248 Kuhnert, P., Overesch, G., 2014. Molecular epidemiology of *Mycoplasma hyopneumoniae* from
249 outbreaks of enzootic pneumonia in domestic pig and the role of wild boar. Vet. Microbiol.
250 174, 261-266.

251 Kuhnert, P., Overesch, G., Belloy, L., 2011. Genotyping of *Mycoplasma hyopneumoniae* in wild
252 boar lung samples. Vet. Microbiol. 152, 191-195.

253 Lin, J.H., Chen, S.P., Yeh, K.S., Weng, C.N., 2006. *Mycoplasma hyorhinis* in Taiwan: diagnosis
254 and isolation of swine pneumonia pathogen. Vet. Microbiol. 115, 111-116.

255 Luehrs, A., 2015. Occurrence of *Mycoplasma hyorhinis* infections in fattening pigs and association
256 with clinical symptoms and pathological lesions of Enzootic Pneumoniae. Thesis of the
257 Vetsuisse Faculty, University of Bern, Bern, Switzerland.

258 Mayor, D., Jores, J., Korczak, B.M., Kuhnert, P., 2008. Multilocus sequence typing (MLST) of
259 *Mycoplasma hyopneumoniae*: A diverse pathogen with limited clonality. Vet. Microbiol.
260 127, 63-72.

261 Mayor, D., Zeeh, F., Frey, J., Kuhnert, P., 2007. Diversity of *Mycoplasma hyopneumoniae* in pig
262 farms revealed by direct molecular typing of clinical material. Vet. Res. 38, 391-398.

263 Nathues, H., Beilage, E.G., Kreienbrock, L., Rosengarten, R., Spergser, J., 2011. RAPD and VNTR
264 analyses demonstrate genotypic heterogeneity of *Mycoplasma hyopneumoniae* isolates from
265 pigs housed in a region with high pig density. Vet. Microbiol. 152, 338-345.

266 Nathues, H., Chang, Y.M., Wieland, B., Rechter, G., Spergser, J., Rosengarten, R., Kreienbrock, L.,
267 Grosse Beilage, E., 2014. Herd-level risk factors for the seropositivity to *Mycoplasma*
268 *hyopneumoniae* and the occurrence of enzootic pneumonia among fattening pigs in areas of
269 endemic infection and high pig density. Transbound. Emerg. Dis. 61, 316-328.

270 Shin, J.H., Joo, H.S., Lee, W.H., Seok, H.B., Calsamig, M., Pijoan, C., Molitor, T.W., 2003.
271 Identification and characterization of cytopathogenic *Mycoplasma hyorhinis* from swine
272 farms with a history of abortions. J. Vet. Med. Sci. 65, 501-509.

273 Stark, K.D., Miserez, R., Siegmann, S., Ochs, H., Infanger, P., Schmidt, J., 2007. A successful
274 national control programme for enzootic respiratory diseases in pigs in Switzerland.
275 Rev.Sci.Tech. 26, 595-606.

276 Switzer, W.P., 1955. Studies on infectious atrophic rhinitis. IV. Characterization of a
277 pleuropneumonia-like organism isolated from the nasal cavities of swine. Am. J. Vet. Res.
278 16, 540-544.

279 Tocqueville, V., Ferre, S., Nguyen, N.H., Kempf, I., Marois-Crehan, C., 2014. Multilocus sequence
280 typing of *Mycoplasma hyorhinis* strains identified by a real-time TaqMan PCR assay. J.
281 Clin. Microbiol. 52, 1664-1671.

282

283

284

285 **Figure legends**

286

287 Fig. 1 Minimal spanning tree of *M. hyorhinis* sequence types (ST) based on allele numbers built in
288 Bionumerics v7.6. A clonal complex relation is defined as a minimum of 4 identical alleles out of
289 the six. Branch length and intensity reflects number of different alleles. The circle size represents
290 the number of isolates belonging to the specific ST which is indicated by the corresponding number.

291

292 Fig. 2 Cluster analysis and strain information of *M. hyorhinis* isolates. A UPGMA tree was built in
293 Bionumerics v7.6 based on the six partial gene sequences used for MLST. The isolate name,
294 sequence type (ST), country and year of isolation, clinic and source are indicated. For the strains
295 analyzed in this study the real-time PCR results for *M. hyopneumoniae* (ABC/REP) are included as
296 well. The bar indicates the genetic distance.

297

298

299 Table 1. Primers used for uniform MLST of *Mycoplasma hyorhinis*

300

301	Gene	Primer	Sequence	seq. (bp) ^a
302	<i>dnaA</i>	dnaA-F2	CAGAAGTCTTAGGTGGTTTTG	459
303		dnaA-R2	TGTGGAATGATCCTTGCCTC	
304				
305	<i>rpoB</i>	rpoB-F2	TCAAGCTGTTCCATTAATTACTAC	509
306		rpoB-R2	GCACTAACTTCTGATCCAATAC	
307				
308	<i>gyrB</i>	gyrB-F2	GATTCTGATGGTTCACATATTAG	358
309		gyrB-R2	GTCTAGGTTTTTTGCATATTTTGC	
310				
311	<i>gltX</i>	gltX-F2	CTGAAAGACTCTCAAAATCACC	448
312		gltX-R2	TTACAAGCCTTTTTGAAATTAGTTC	
313				
314	<i>adk</i>	adk-F2	CGATGGCATCTAATTCTTTTAAAG	437
315		adk-R2	TACTCAGGCAAAGTTTTTAGAAC	
316				
317	<i>gmk</i>	gmk-F2	GCGCCTGTTTCTGTTAATATTG	452
318		gmk-R2	AAGAGACAAAAGACCTAATGAAG	

319 ^a length of the informative sequence, i.e. of the PCR product without primer sequences

320

321

Table 1. Primers used for uniform MLST of *Mycoplasma hyorhinis*

Gene	Primer	Sequence	seq. (bp) ^a
<i>dnaA</i>	dnaA-F2	CAGAAAGTCTTAGGTGGTTTTG	459
	dnaA-R2	TGTGGAATGATCCTTGCCTC	
<i>rpoB</i>	rpoB-F2	TCAAGCTGTTCCATTAATTACTAC	509
	rpoB-R2	GCACTAACTTCTGATCCAATAC	
<i>gyrB</i>	gyrB-F2	GATTCTGATGGTTCACATATTAG	358
	gyrB-R2	GTCTAGGTTTTTTGCATATTTTGC	
<i>gltX</i>	gltX-F2	CTGAAAGACTCTCAAAATCACC	448
	gltX-R2	TTACAAGCCTTTTTGAAATTAGTTC	
<i>adk</i>	adk-F2	CGATGGCATCTAATTCTTTTAAAG	437
	adk-R2	TACTCAGGCAAAGTTTTAGAAC	
<i>gmk</i>	gmk-F2	GCGCCTGTTTCTGTTAATATTG	452
	gmk-R2	AAGAGACAAAAGACCTAATGAAG	

^a length of the informative sequence, i.e. of the PCR product without primer sequences

Fig. 1 Minimal spanning tree.