

A FOUR MOMENTS THEOREM FOR GAMMA LIMITS ON A POISSON CHAOS

TOBIAS FISSLER AND CHRISTOPH THÄLE

ABSTRACT. This paper deals with sequences of random variables belonging to a fixed chaos of order q generated by a Poisson random measure on a Polish space. The problem is investigated whether convergence of the third and fourth moment of such a suitably normalized sequence to the third and fourth moment of a centred Gamma law implies convergence in distribution of the involved random variables. A positive answer is obtained for $q = 2$ and $q = 4$. The proof of this four moments theorem is based on a number of new estimates for contraction norms. Applications concern homogeneous sums and U -statistics on the Poisson space.

1. INTRODUCTION

Probabilistic limit theorems for sequences of multiple stochastic integrals have found considerable attention during the last decade. One of the most remarkable results in this direction is the fourth moment theorem of Nualart and Peccati obtained in the seminal paper [12]. It asserts that a sequence of suitably normalized multiple stochastic integrals of order $q \geq 1$ with respect to a Gaussian random measure on a Polish space satisfies a central limit theorem if and only if the sequence of their fourth moments converges to 3, the fourth moment of a standard Gaussian distribution. This drastic simplification of the method of moments has stimulated a large number of applications, for example to Gaussian random processes or fields, mathematical statistics, random matrices or random polynomials (we refer the reader to the monograph [10] and also to the constantly updated webpage <https://sites.google.com/site/malliavinstein/home> for further details and references).

Besides the fourth moment theorem mentioned above, there is also a ‘non-central’ version dealing with the approximation of a sequence of multiple stochastic integrals by a Gamma-distributed random variable, cf. [9]. Again, the result is a drastic simplification of the method of moments as it delivers convergence in distribution if and only if a certain linear combination of the third and the fourth moment of the involved random variables converges to the corresponding expression for Gamma random variables. In view of normalization conditions we see that in fact the first four moments of the random variables are involved, which gives rise to the name ‘four moments theorem’ for such a result. To simplify the terminology, we will also speak about a four moments theorem in the case of normal approximation.

The present paper asks whether a similar non-central limit theorem is available for sequences of multiple stochastic integrals with respect to a Poisson random measure on a Polish space.

2010 *Mathematics Subject Classification*. Primary: 60F05, 60H05, 60G57; Secondary: 60G55, 62E20.

Key words and phrases. Contraction, four moments theorem, Gamma distribution, homogeneous sums, non-central limit theorem, Poisson chaos, Poisson random measure, universality, U -statistics.

The first author has been supported by the Swiss National Science Foundation (SNF) via grant 152609. The second author has been supported by the German Research Foundation (DFG) via SFB-TR 12.

In this set-up, a central four moments theorem has been derived by Lachièze-Rey and Peccati in [6] under an additional sign condition (see also [5]), which, on the Poisson space, seems to be unavoidable. While Gamma approximation on the Poisson space in the spirit of the Malliavin-Stein method has been dealt with in [17], the problem of a four moments theorem similar to that for Gaussian multiple stochastic integrals mentioned above remained open in general. The main result of our paper, Theorem 3.5, delivers a four moments theorem for sequences of Poisson stochastic integrals of order $q = 2$ and $q = 4$. For this reason, the present work can be seen as a natural continuation of [17], where the case $q = 2$ has already been settled under additional assumptions, which we are able to overcome. The proof of our four moments theorem relies on a couple of new estimates for norms of so-called contraction kernels and the combinatorially involved multiplication formula for stochastic integrals on the Poisson space. It is precisely this combinatorial complexity which forces that our proof yields a positive result only for sequences of Poisson stochastic integrals of order $q = 2$ and $q = 4$. However, all intermediate steps in our proof will be formulated for general $q \geq 2$ to make as transparent as possible and to highlight, in which argument the restrictive condition on the order of the integrals arises. The main difference between the central and the non-central version of the four moments theorem is that in the non-central case one has to deal with a linear combination of the third and the fourth moment of the stochastic integrals, while the central case only requires an analysis of the fourth moment. Even under additional conditions on the integrands, this leads to difficulties, which we can overcome only for $q = 2$ and $q = 4$. We have to leave it as an open problem for future research to extend our result to arbitrary q by other methods.

The main result of our paper is applied to a universality question for homogeneous sums on a Poisson chaos as well as to a non-central analogue of de Jong's theorem for completely degenerate U -statistics of order two and four. This partially complements the results for Gamma and normal approximation obtained in [5, 17] and [19]. We emphasize in this context that limit theorems for non-linear functionals of Poisson random measures have recently found numerous applications especially in geometric probability or stochastic geometry [5, 6, 7, 8, 17, 22, 23] and in the theory of Lévy processes [5, 7, 14, 18].

Our paper is structured as follows. In Section 2, we introduce and collect necessary background material. To contrast our results with those available for Gaussian multiple stochastic integrals, we shall present them in the context of completely random measures, which captures both settings. Our main results are the content of Section 3, while Section 4 contains applications to homogeneous sums and U -statistics. The proof of Theorem 3.5 is presented in the final Section 5.

2. PRELIMINARIES

In this section, we introduce the basic definitions, mainly related to Poisson stochastic integrals. For further details and background material we refer the reader to the monograph [16] as well as to the papers [13, 14].

2.1. Completely random measures. Without loss of generality, we assume that all objects are defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let \mathcal{Z} denote a Polish space with Borel σ -field \mathcal{L} , which is equipped with a non-atomic σ -finite measure μ . We define the class $\mathcal{L}_\mu = \{B \in \mathcal{L} : \mu(B) < \infty\}$ and let $\varphi = \{\varphi(B) : B \in \mathcal{L}_\mu\}$ indicate a *completely random measure* on $(\mathcal{Z}, \mathcal{L})$ with control measure μ . That is, φ is a set of random variables such that

- (i) for every collection of pairwise disjoint elements $B_1, \dots, B_n \in \mathcal{Z}_\mu$, the random variables $\varphi(B_1), \dots, \varphi(B_n)$ are independent;
- (ii) for every $B, C \in \mathcal{Z}_\mu$, one has the identity $\mathbb{E}[\varphi(B)\varphi(C)] = \mu(B \cap C)$.

If $\mathbb{E}[\varphi(B)] = 0$ and $\varphi(B) \in L^2(\mathbb{P})$ (i.e., $\varphi(B)$ is square-integrable with respect to \mathbb{P}) for every $B \in \mathcal{Z}_\mu$, then the mapping $\mathcal{Z}_\mu \rightarrow L^2(\mathbb{P}), B \mapsto \varphi(B)$, is σ -additive in the sense that for every sequence $(B_n)_{n \geq 1}$ of pairwise disjoint elements of \mathcal{Z}_μ , one has that

$$\varphi \left(\bigcup_{n=1}^{\infty} B_n \right) = \sum_{n=1}^{\infty} \varphi(B_n) \quad \mathbb{P}\text{-a.s.}, \quad (2.1)$$

where the right-hand side converges in $L^2(\mathbb{P})$. By $\sigma(\varphi)$ we denote the σ -field generated by φ . In this paper, we shall deal with two special and prominent instances of completely random measures, namely a centred Gaussian and a compensated Poisson measure.

- (a) A *centred Gaussian measure* with control measure μ is denoted by G and is a completely random measure such that the elements of G are jointly Gaussian and centred.
- (b) A *compensated Poisson measure* with control measure μ is indicated by $\hat{\eta}$ and is a completely random measure such that for every $B \in \mathcal{Z}_\mu$, $\hat{\eta}(B) \stackrel{d}{=} \eta(B) - \mu(B)$, where $\eta(B)$ is a Poisson random variable with mean $\mu(B)$.

By definition, both G and $\hat{\eta}$ are centred families in $L^2(\mathbb{P})$, implying that (2.1) is satisfied. Moreover, for \mathbb{P} -almost every $\omega \in \Omega$, $\hat{\eta}(\cdot, \omega)$ is a signed measure on $(\mathcal{Z}, \mathcal{Z})$, while G does not satisfy this property, cf. [16, Example 5.1.7 (iii)].

2.2. L^2 -spaces. Let $q \geq 1$ be an integer. We shall use the shorthand notation $L^2(\mu^q)$ for the space $L^2(\mathcal{Z}^q, \mathcal{Z}^q, \mu^q)$ of (deterministic) functions that are square-integrable with respect to μ^q . $L_s^2(\mu^q)$ stands for the subspace of $L^2(\mu^q)$ consisting of symmetric functions, i.e., functions that are μ^q -a.e. invariant under permutations of their arguments. For $f, g \in L^2(\mu^q)$ we define the scalar product $\langle f, g \rangle_{L^2(\mu^q)} = \int_{\mathcal{Z}^q} fg \, d\mu^q$ and the norm $\|f\|_{L^2(\mu^q)} = \langle f, f \rangle_{L^2(\mu^q)}^{1/2}$. If there is no risk of confusion, we suppress in what follows the dependency on q and μ , and merely write $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively. Moreover, let $L^2(\sigma(\varphi), \mathbb{P})$ denote the space of all square-integrable functionals of φ , where φ is either a Poisson measure $\hat{\eta}$ or a Gaussian measure G . If $F \in L^2(\sigma(\varphi), \mathbb{P})$, we shall sometimes write $F = F(\varphi)$ in order to underpin the dependency of F on φ . As a convention, we shall use lower case variables for elements of $L^2(\mu^q)$ and capitals for elements of $L^2(\sigma(\varphi), \mathbb{P})$. Finally, we introduce the space $L^2(\mathbb{P}, L^2(\mu)) = L^2(\Omega \times \mathcal{Z}, \mathcal{F} \otimes \mathcal{Z}, \mathbb{P} \otimes \mu)$ as the space of all jointly square-integrable measurable mappings $u: \Omega \times \mathcal{Z} \rightarrow \mathbb{R}$. If $u, v \in L^2(\mathbb{P}, L^2(\mu))$, their scalar product is defined as $\langle u, v \rangle_{L^2(\mathbb{P}, L^2(\mu))} = \int_{\Omega} \int_{\mathcal{Z}} u(\omega, z)v(\omega, z)\mu(dz)\mathbb{P}(d\omega)$ and we denote by $\|\cdot\|_{L^2(\mathbb{P}, L^2(\mu))}$ the norm induced by it.

2.3. Multiple stochastic integrals. Let $\varphi = \hat{\eta}$ or $\varphi = G$. For every integer $q \geq 1$ we denote the *multiple stochastic integral* of order q with respect to φ by I_q^φ . It is a mapping $I_q^\varphi: L_s^2(\mu^q) \rightarrow L^2(\sigma(\varphi), \mathbb{P})$, which is linear and continuous. Additionally, for $f \in L_s^2(\mu^q)$, the random variable $I_q^\varphi(f)$ is centred. Moreover, the multiple stochastic integral satisfies the *Itô isometry*

$$\mathbb{E}[I_p^\varphi(f)I_q^\varphi(g)] = \begin{cases} 0 & \text{if } q \neq p \\ q! \langle f, g \rangle_{L_s^2(\mu^q)} & \text{if } q = p \end{cases} \quad (2.2)$$

for any integers $p, q \geq 1$ and $f \in L_s^2(\mu^p)$, $g \in L_s^2(\mu^q)$. For general $f \in L^2(\mu^q)$, we put $I_q^\varphi(f) = I_q^\varphi(\tilde{f})$, where

$$\tilde{f}(z_1, \dots, z_q) = \frac{1}{q!} \sum_{\pi \in \Pi_q} f(z_{\pi(1)}, \dots, z_{\pi(q)})$$

is the canonical symmetrization of f , and Π_q is the group of all $q!$ permutations π of the set $\{1, \dots, q\}$. We emphasize that due to Jensen's inequality and the convexity of norms, we have the inequality $\|\tilde{f}\| \leq \|f\|$. As a convention, we set $I_0^\varphi: \mathbb{R} \rightarrow \mathbb{R}$ equal to the identity map on \mathbb{R} .

Since this article is mostly concerned with Poisson integrals, we shall write I_q instead of $I_q^{\hat{\eta}}$.

2.4. Chaos decomposition. The Itô isometry in (2.2) formalizes an orthogonality relation between multiple stochastic integrals of different order. This induces the following so-called *chaos decomposition* (see [13]):

$$L^2(\sigma(\varphi), \mathbb{P}) = \bigoplus_{q=0}^{\infty} W_q^\varphi, \quad (2.3)$$

where $W_0^\varphi = \mathbb{R}$ and $W_q^\varphi = \{I_q^\varphi(f) : f \in L_s^2(\mu^q)\}$ for $\varphi = \hat{\eta}$ or $\varphi = G$, $q \geq 1$. Depending on the choice of φ , we shall often use the terms *Poisson chaos* and *Gaussian chaos of order q* for W_q^φ , respectively.

A consequence of (2.3) is that any $F \in L^2(\sigma(\varphi), \mathbb{P})$, with $\varphi = \hat{\eta}$ or $\varphi = G$, admits a chaotic decomposition

$$F = \mathbb{E}[F] + \sum_{q=1}^{\infty} I_q^\varphi(f_q),$$

where the kernels $f_q \in L_s^2(\mu^q)$ are unique μ^q -a.e. and the series converges in $L^2(\mathbb{P})$.

2.5. Contractions. Fix integers $p, q \geq 1$ and functions $f \in L_s^2(\mu^p)$, $g \in L_s^2(\mu^q)$. For any $r \in \{0, \dots, p \wedge q\}$ and $\ell \in \{1, \dots, r\}$ we define the *contraction* $f \star_r^\ell g: \mathcal{Z}^{p+q-r-\ell} \rightarrow \mathbb{R}$ which acts on the tensor product $f \otimes g$ and reduces the number of variables from $p+q$ to $p+q-r-\ell$ in the following way: r variables are identified and among these, ℓ are integrated out with respect to μ . More formally,

$$\begin{aligned} & f \star_r^\ell g(\gamma_1, \dots, \gamma_{r-\ell}, t_1, \dots, t_{p-r}, s_1, \dots, s_{q-r}) \\ &= \int_{\mathcal{Z}^\ell} f(z_1, \dots, z_\ell, \gamma_1, \dots, \gamma_{r-\ell}, t_1, \dots, t_{p-r}) \\ & \quad \times g(z_1, \dots, z_\ell, \gamma_1, \dots, \gamma_{r-\ell}, s_1, \dots, s_{q-r}) \mu^\ell(d(z_1 \dots z_\ell)), \end{aligned}$$

and for $\ell = 0$ we put

$$\begin{aligned} & f \star_r^0 g(\gamma_1, \dots, \gamma_r, t_1, \dots, t_{p-r}, s_1, \dots, s_{q-r}) \\ &= f(\gamma_1, \dots, \gamma_r, t_1, \dots, t_{p-r}) g(\gamma_1, \dots, \gamma_r, s_1, \dots, s_{q-r}). \end{aligned}$$

Note that even if f and g are symmetric, the contraction $f \star_r^\ell g$ is not necessarily symmetric. We denote the canonical symmetrization by

$$f \tilde{\star}_r^\ell g(z_1, \dots, z_{p+q-r-\ell}) = \frac{1}{(p+q-r-\ell)!} \sum_{\pi \in \Pi_{p+q-r-\ell}} f \star_r^\ell g(z_{\pi(1)}, \dots, z_{\pi(p+q-r-\ell)}).$$

We also emphasize that for $f \in L_s^2(\mu^p)$ and $g \in L_s^2(\mu^q)$, the contraction $f \star_r^\ell g$ is neither necessarily well-defined nor necessarily an element of $L^2(\mu^{p+q-r-\ell})$. At least, by using the Cauchy-Schwarz inequality, we can deduce that $f \star_r^\ell g \in L^2(\mu^{p+q-2r})$ for any $r \in \{0, \dots, p \wedge q\}$. For this reason and to circumvent any complications in the calculations, we make the following technical assumptions.

2.6. Technical assumptions (A). We use the same set of technical assumptions as in [6, 14, 17]. For a detailed explanation of the conditions and their consequences, we refer to these works.

For a sequence $F_n = I_q(f_n)$ of multiple integrals of fixed order $q \geq 1$ with $f_n \in L^2(\mu_n^q)$ for every $n \geq 1$ (we allow the non-atomic and σ -finite measure to vary with n), we assume that the following three technical conditions are satisfied:

- (a) for any $r \in \{1, \dots, q\}$, the contraction $f_n \star_q^{q-r} f_n$ is an element of $L^2(\mu_n^r)$;
- (b) for any $r \in \{1, \dots, q\}$, $\ell \in \{1, \dots, r\}$ and $(z_1, \dots, z_{2q-r-\ell}) \in \mathcal{Z}^{2q-r-\ell}$, we have that $(|f_n| \star_r^\ell |f_n|)(z_1, \dots, z_{2q-r-\ell})$ is well-defined and finite;
- (c) for any $k \in \{0, \dots, 2(q-1)\}$ and any r and ℓ satisfying $k = 2(q-1) - r - \ell$, we have that

$$\int_{\mathcal{Z}} \sqrt{\int_{\mathcal{Z}^k} (f_n(z, \cdot) \star_r^\ell f_n(z, \cdot))^2 d\mu_n^k \mu_n(dz)} < \infty.$$

2.7. Multiplication formula. A very convenient property of multiple stochastic integrals is that one can express the product of two such integrals as a linear combination of multiple integrals of contraction kernels. More precisely, we have the following multiplication formula for Poisson integrals, which is taken from [16, Proposition 6.5.1].

Lemma 2.1 (Multiplication formula for Poisson integrals). *Let $f \in L_s^2(\mu^p)$ and $g \in L_s^2(\mu^q)$, $p, q \geq 1$. Suppose that $f \star_r^\ell g \in L^2(\mu^{p+q-r-\ell})$ for every $r \in \{0, \dots, p \wedge q\}$ and every $\ell \in \{0, \dots, r\}$. Then*

$$I_p(f)I_q(g) = \sum_{r=0}^{p \wedge q} r! \binom{p}{r} \binom{q}{r} \sum_{\ell=0}^r I_{p+q-r-\ell}(f \tilde{\star}_r^\ell g). \quad (2.4)$$

We remark that if a kernel $f \in L_s^2(\mu^q)$ satisfies the technical assumptions (A), the assumptions of Lemma 2.1 are automatically satisfied if $g = f$, implying that $I_q(f)^2 \in L^2(\sigma(\hat{\eta}), \mathbb{P})$. To simplify our notation, for $f \in L_s^2(\mu^q)$ we put $G_0^q f = q! \|f\|^2$ and

$$G_p^q f = \sum_{r=0}^q \sum_{\ell=0}^r \mathbf{1}(2q-r-\ell=p) r! \binom{q}{r}^2 \binom{r}{\ell} f \tilde{\star}_r^\ell f \quad (2.5)$$

for $p \in \{1, \dots, 2q\}$. In other words, the operator G_p^q turns a function of q variables into a function of p variables. We can now re-write (2.4) in a simplified form as

$$I_q(f)^2 = \sum_{p=0}^{2q} I_p(G_p^q f)$$

with $I_0(G_0^q f) = G_0^q f = q! \|f\|^2$.

The multiplication formula paves the way for the computation of moments of multiple stochastic integrals. In particular, we have the following expressions for the third and the fourth moment of a multiple Poisson integral.

Lemma 2.2 (Third and fourth moment of Poisson integrals). *Fix an integer $q \geq 1$. Let $f \in L_s^2(\mu^q)$ such that the technical assumptions (A) are satisfied. Then $I_q(f) \in L^4(\mathbb{P})$. Moreover, we have that*

$$\mathbb{E}[I_q(f)^3] = q! \sum_{r=0}^q \sum_{\ell=0}^r \mathbf{1}(q = r + \ell) r! \binom{q}{r}^2 \binom{r}{\ell} \langle f \tilde{\star}_r^\ell f, f \rangle, \quad (2.6)$$

$$\mathbb{E}[I_q(f)^4] = \sum_{p=0}^{2q} p! \|G_p^q f\|^2. \quad (2.7)$$

Proof. The technical assumptions (A) ensure that all symmetrized contraction kernels $f \tilde{\star}_r^\ell f$ appearing in (2.6) and (2.7) are elements of $L^2(\mu^{2q-r-\ell})$, which implies that the third and the fourth moment of $I_q(f)$ are finite. The explicit formulae in (2.6) and (2.7) follow directly from the isometry property (2.2) and the multiplication formula (2.4). \square

Remark 2.3. Note that for even $q \geq 2$, (2.6) reduces to

$$\mathbb{E}[I_q(f)^3] = q! \sum_{r=q/2}^q r! \binom{q}{r}^2 \binom{r}{q-r} \langle f \tilde{\star}_r^{q-r} f, f \rangle. \quad (2.8)$$

Remark 2.4. There is also a multiplication formula for the Gaussian case. It reads

$$I_p^G(f) I_q^G(g) = \sum_{r=0}^{p \wedge q} r! \binom{p}{r} \binom{q}{r} I_{p+q-2r}^G(f \tilde{\star}_r^r g),$$

where $p, q \geq 1$ and $f \in L_s^2(\mu^p)$, $g \in L_s^2(\mu^q)$, see [16, Proposition 6.4.1]. As a consequence, we see that the third and fourth moment of a Gaussian multiple integral have a more compact form compared to the Poisson case. Indeed, for an integer $q \geq 1$ and $f \in L_s^2(\mu^q)$, one has that

$$\mathbb{E}[I_q^G(f)^3] = \frac{(q!)^3}{(q/2!)^2} \langle f \tilde{\star}_{q/2}^{q/2} f, f \rangle \mathbf{1}(q \text{ is even}), \quad (2.9)$$

$$\mathbb{E}[I_q^G(f)^4] = \sum_{r=0}^q (r!)^2 \binom{q}{r}^4 (2q - 2r)! \|f \tilde{\star}_r^r f\|^2.$$

In particular, the third moment of a Gaussian integral of odd order vanishes, while this is in general not the case for a Poisson integral.

3. FOUR MOMENTS THEOREMS

This section contains the main results of our paper, namely a four moments theorem for Gamma approximation on a Poisson chaos of fixed order. To allow for an easier comparison with the existing literature, we first recall known results on a Gaussian chaos and also a version of the four moments theorem for normal approximation on a Poisson chaos.

3.1. Four moments theorems on a Gaussian chaos. The classical method of moments yields a central limit theorem for a normalized sequence of random variables under the condition that all moments converge to those of the standard Gaussian distribution. The four moments theorem on a Gaussian chaos is a drastical simplification of the method of moments as it provides a central limit theorem for a sequence of normalized Gaussian multiple stochastic integrals under the much weaker condition that only the fourth moment converges

to 3 (which is the fourth moment of the standard Gaussian distribution). Alternatively, this statement can be re-formulated in terms of the convergence of norms of contractions. In what follows we write $X \sim \mathcal{L}$ if a random variable X has distribution \mathcal{L} .

Theorem 3.1 (see Theorem 1 in [12]). *Fix an integer $q \geq 2$ and let $\{f_n: n \geq 1\} \subset L_s^2(\mu^q)$ be such that*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q^G(f_n)^2] = 1.$$

Further, let $N \sim \mathcal{N}(0, 1)$ be a standard Gaussian random variable. Then the following three assertions are equivalent:

- (i) *As $n \rightarrow \infty$, the sequence $\{I_q^G(f_n): n \geq 1\}$ converges in distribution to N ;*
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[I_q^G(f_n)^4] = 3$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^r f_n\| = 0$ for every $r \in \{1, \dots, q-1\}$.

In the subsequent work [9], the authors have shown a ‘non-central’ version of Theorem 3.1 where the limiting distribution is a centred Gamma distribution. To state the result properly, let us recall the formal definition of the latter limit law.

Definition 3.2 (Centred Gamma distribution). A random variable Y has a centred Gamma distribution $\bar{\Gamma}_\nu$ with parameter $\nu > 0$, if

$$Y \stackrel{d}{=} 2X - \nu,$$

where X has the usual Gamma law with mean and variance both equal to $\nu/2$ and where $\stackrel{d}{=}$ stands for equality in distribution. The probability density of $\bar{\Gamma}_\nu$ is given by

$$g_\nu(x) = \frac{2^{-\nu/2}}{\Gamma(\nu/2)} (x + \nu)^{\nu/2-1} e^{-(x+\nu)/2} \mathbf{1}(x > -\nu),$$

and the the first four moments of Y are

$$\mathbb{E}[Y] = 0, \quad \mathbb{E}[Y^2] = 2\nu, \quad \mathbb{E}[Y^3] = 8\nu, \quad \mathbb{E}[Y^4] = 12\nu^2 + 48\nu.$$

We are now in the position to re-phrase the following non-central analogue of Theorem 3.1.

Theorem 3.3 (see Theorem 1.2 in [9]). *Let $\nu > 0$ and fix an even integer $q \geq 2$. Let $\{f_n: n \geq 1\} \subset L_s^2(\mu^q)$ be such that*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q^G(f_n)^2] = 2\nu.$$

Further, let $Y \sim \bar{\Gamma}_\nu$ be a centred Gamma-distributed random variable with parameter ν . Then the following three assertions are equivalent:

- (i) *As $n \rightarrow \infty$, the sequence $\{I_q^G(f_n): n \geq 1\}$ converges in distribution to Y ;*
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[I_q^G(f_n)^4] - 12\mathbb{E}[I_q^G(f_n)^3] = 12\nu^2 - 48\nu$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^r f_n\| = 0$ for every $r \in \{1, \dots, q-1\} \setminus \{q/2\}$, and

$$\lim_{n \rightarrow \infty} \|f_n \star_{q/2}^{q/2} f_n - c_q f_n\| = 0$$
 with $c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}$.

It is a characterizing feature of the centred Gamma-distribution that the so-called ‘middle-contraction’ $f_n \star_{q/2}^{q/2} f_n$ plays a special role in condition (iii). The fact that the middle-contraction does not vanish goes hand in hand with the appearance of the third moment in condition (ii), recall (2.9).

3.2. Four moments theorems on a Poisson chaos. We now turn to four moments theorems on a Poisson chaos of fixed order $q \geq 2$. To this end, let, for each $n \geq 1$, μ_n be a σ -finite non-atomic measure on $(\mathcal{Z}, \mathcal{Z})$ and denote by $\hat{\eta}_n$ a compensated Poisson random measure with control μ_n . Further let $\{f_n : n \geq 1\}$ be a sequence of symmetric function such that f_n is square-integrable with respect to μ_n^q for each $n \geq 1$. In this set-up, $\|f_n\|$ denotes the norm of f_n with respect to μ_n^q , and $f_n \star_r^\ell f_n$ stands for the contraction taken with respect to μ_n . Finally, define $F_n = I_q(f_n)$, where for each n the stochastic integral is with respect to $\hat{\eta}_n$. As in the Gaussian case discussed in the previous section, we start with the case of a standard normal limiting distribution.

Theorem 3.4 (see Theorem 3.12 in [6]). *Let $\{\mu_n : n \geq 1\}$ be a sequence of σ -finite and non-atomic measures such that $\lim_{n \rightarrow \infty} \mu_n(\mathcal{Z}) = \infty$ and fix $q \geq 2$. Let $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence such that for each $n \geq 1$ either $f_n \geq 0$ or $f_n \leq 0$. Suppose that the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = 1 \quad (3.1)$$

are satisfied. Further, suppose that $\{I_q(f_n)^4 : n \geq 1\}$ is uniformly integrable and let $N \sim \mathcal{N}(0, 1)$ be a standard Gaussian random variable. Then the following three assertions are equivalent:

- (i) As $n \rightarrow \infty$, the sequence $\{I_q(f_n) : n \geq 1\}$ converges in distribution to N ;
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^4] = 3$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0$ for all $r \in \{1, \dots, q\}$ and $\ell \in \{1, \dots, r \wedge (q - 1)\}$, and $\lim_{n \rightarrow \infty} \|f_n\|_{L^4(\mu_n^q)} = 0$.

Let us comment on the differences between Theorem 3.1 and Theorem 3.4.

- (1) In the Poisson case one has to ensure that the involved control measures are infinite measures, at least in the limit, as $n \rightarrow \infty$. The reason for this is that otherwise, the normalization (3.1) and the condition that $\lim_{n \rightarrow \infty} \|f_n\|_{L^4(\mu_n^q)} = 0$ are mutually exclusive, see also the remark after Assumption N in [15] for a brief discussion of this problem.
- (2) One has to assume that the functions f_n have a constant sign, that is for each $n \geq 1$ either $f_n \geq 0$ or $f_n \leq 0$. The reason for this is that in the Poisson case, besides of the contraction norms $\|f_n \star_r^\ell f_n\|$, also scalar products of the form $\langle f_n \star_{r_1}^{\ell_1} f_n, f_n \star_{r_2}^{\ell_2} f_n \rangle$ enter the expression of the fourth moments $\mathbb{E}[I_q(f_n)^4]$. The sign condition then allows to control the signs of these scalar products, which rules out cancellation effects.
- (3) In the Poisson case, one also has to assume that the sequence $\{I_q(f_n)^4 : n \geq 1\}$ is uniformly integrable, while in the Gaussian case, this condition is automatically fulfilled thanks to the hypercontractivity property of Gaussian integrals (see e.g. [10, Theorem 2.7.2]). This is needed to ensure that the convergence in distribution of $I_q(f_n)$ to N implies the convergence of the first four moments.

For general $q \geq 2$ and general sequences $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ there is no version of a four moments theorem on a Poisson chaos relaxing one of the conditions discussed above. However, for $q = 2$ the sign condition is not necessary as shown by Theorem 2 in [15]. Moreover, for general $q \geq 2$ and if the sequence $\{f_n : n \geq 1\}$ is *tamed* (see Definition 4.2 below), Theorem 3.2 in [19] provides a four moments theorem without a sign condition. In this case, also

condition (iii) can be relaxed by assuming – besides the condition on the L^4 -norm of f_n – only that $\lim_{n \rightarrow \infty} \|f_n \star_r^r f_n\| = 0$ for all $r \in \{1, \dots, q-1\}$.

After having discussed the four moments theorem for normal approximation on the Poisson space, we now turn to the main result of the present work, namely a version of Theorem 3.3 for Poisson integrals of order $q = 2$ and $q = 4$. The reason for this rather restrictive condition on the order of the involved integrals will be discussed below.

Theorem 3.5 (Four moments theorem for Poisson integrals). *Fix $\nu > 0$. Let $q \geq 2$ be even and $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence satisfying the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = 2\nu.$$

Furthermore, let the sequence $\{I_q(f_n)^4: n \geq 1\}$ be uniformly integrable and let $Y \sim \bar{\Gamma}_\nu$ be a random variable following a centred Gamma distribution with parameter ν . If one of the conditions

- (a) $q = 2$ and $\lim_{n \rightarrow \infty} \|f_n^2\| = 0$,
- (b) $q = 4$ and $f_n \leq 0$ for all $n \geq 1$

is satisfied, then the following three assertions are equivalent:

- (i) As $n \rightarrow \infty$, the sequence $\{I_q(f_n): n \geq 1\}$ converges in distribution to Y ;
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3] = 12\nu^2 - 48\nu$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0$ for all $r \in \{1, \dots, q\}$ and $\ell \in \{1, \dots, r \wedge (q-1)\}$ such that $(r, \ell) \neq (q/2, q/2)$, $\lim_{n \rightarrow \infty} \|f_n\|_{L^4(\mu_n^q)} = 0$, and $\lim_{n \rightarrow \infty} \|f_n \star_{q/2}^{q/2} f_n - c_q f_n\| = 0$ with $c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}$.

Remark 3.6. Under condition (a), Theorem 3.5 is a version of Proposition 2.9 in [17]. However, in that paper one has to assume that for each $n \geq 1$ the reference measure μ_n is finite. As discussed earlier in this section, this is a quite restrictive assumption. We provide a proof which circumvents this technicality.

The implication (i) \implies (ii) of Theorem 3.5 is a direct consequence of the uniform integrability assumption. That (iii) implies (i) follows from a generalization of Theorem 2.6 in [17] stated as Proposition 5.1 below. Showing the implication (ii) \implies (iii) is the main part of the proof. While the proof of the corresponding implication in Theorem 3.4 is rather straight forward and works for arbitrary $q \geq 2$, the proof here is based on a couple of new estimates and arguments. They are of independent interest and might also be helpful beyond the context of the present paper. In sharp contrast to Theorem 3.4, our arguments show that the ‘usual’ technique (relying on the multiplication formula for Poisson integrals similar as in the proofs of Theorems 3.1, 3.3 or 3.4) for proving the implication (ii) \implies (iii) only works in case that $q = 2$ and $q = 4$ and cannot be improved. The main reason for this is the involved combinatorial structure on a Poisson chaos implied by the multiplication formula (2.4). The proof of Theorem 3.5 is the content of Section 5 below.

Theorem 3.5 has a counterpart in a free probability setting, see [1]. Here, one studies the approximation of the law of a sequence of elements belonging to a fixed chaos of order $q \geq 1$ of the so-called *free* Poisson algebra by the Marchenko-Pastur law (also called free Poisson law). It is interesting to see that in this case, the proof works for arbitrary $q \geq 1$ and does not need a sign condition on the kernels. This is explained by the relatively simple combinatorial

structure on a free Poisson chaos, which is inherited from the free multiplication formula in which all combinatorial coefficients are equal to one. This causes that the expressions for the third and fourth moment are much simpler compared to the classical set-up of the present paper and implies that a proof of the corresponding free four moments theorem works in full generality.

Comparing Theorem 3.4 and Theorem 3.5, it is natural to ask whether there exists a version of Theorem 3.5 dealing with a sequence of non-negative kernels. Indeed, Corollary 3.8 below provides such a version, but it deals with a different limiting law, namely what we call a centred reflected Gamma distribution. In case of a limiting Gaussian law, this phenomenon is not visible, since a Gaussian law is symmetric, see also the discussion in Remark 5.10.

Definition 3.7 (Centred reflected Gamma distribution). A random variable Y has a centred reflected Gamma distribution $\widehat{\Gamma}_\nu$ with parameter $\nu > 0$, if $-Y \sim \overline{\Gamma}_\nu$.

Note that if $Y \sim \widehat{\Gamma}_\nu$ follows a centred reflected Gamma distribution with parameter ν , the first four moments of Y are given by

$$\mathbb{E}[Y] = 0, \quad \mathbb{E}[Y^2] = 2\nu, \quad \mathbb{E}[Y^3] = -8\nu, \quad \mathbb{E}[Y^4] = 12\nu^2 + 48\nu.$$

Moreover, while the centred Gamma distribution has support $[-\nu, \infty)$, the centred reflected Gamma distribution is supported on $(-\infty, \nu]$. The next result is an immediate consequence of Theorem 3.5 and the definition of $\widehat{\Gamma}_\nu$.

Corollary 3.8 (Four moments theorem for Poisson integrals with non-negative kernels). *Fix $\nu > 0$. Let $q \geq 2$ be an even integer and $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence of kernels satisfying the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = 2\nu.$$

Let the sequence $\{I_q^4(f_n) : n \geq 1\}$ be uniformly integrable and suppose that $Y \sim \widehat{\Gamma}_\nu$ is a random variable having a centred reflected Gamma distribution with parameter ν . If one of the conditions

- (a) $q = 2$ and $\lim_{n \rightarrow \infty} \|f_n^2\| = 0$,
- (b) $q = 4$ and $f_n \geq 0$ for all $n \geq 1$

is satisfied, then the following three assertions are equivalent:

- (i) *As $n \rightarrow \infty$, the sequence $\{I_q(f_n) : n \geq 1\}$ converges in distribution to Y ;*
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^4] + 12\mathbb{E}[I_q(f_n)^3] = 12\nu^2 - 48\nu$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0$ for all $r \in \{1, \dots, q\}$, $\ell \in \{1, \dots, r \wedge (q-1)\}$ such that $(r, \ell) \neq (q/2, q/2)$, $\lim_{n \rightarrow \infty} \|f_n^2\| = 0$, and $\lim_{n \rightarrow \infty} \|f_n \widetilde{\star}_{q/2}^{q/2} f_n + c_q f_n\| = 0$ with $c_q = \frac{4}{(q/2)!(q/2)^2}$.

Remark 3.9. We emphasize that one could derive our main result, Theorem 3.5, also for the two-parametric centred Gamma distribution $\overline{\Gamma}_{a,\lambda}$, $a, \lambda > 0$, with probability density

$$h_{a,\lambda}(x) = \frac{\lambda^a}{\Gamma(a)} \left(x + \frac{a}{\lambda}\right)^{a-1} e^{-(\lambda x + a)} \mathbf{1}(x > -\frac{a}{\lambda}).$$

The one-parametric centred Gamma distribution $\overline{\Gamma}_\nu$ then arises by putting $a = \frac{\nu}{2}$ and $\lambda = \frac{1}{2}$. In order to allow for a better comparison with the existing literature [9, 17] and to keep the presentation transparent, we have decided to restrict to the one-parametric case.

4. APPLICATION TO HOMOGENEOUS SUMS AND U -STATISTICS

4.1. Homogeneous sums. According to [19] a universality result is a ‘mathematical statement implying that the asymptotic behaviour of a large random system does not depend on the distribution of its components’. Such results are at the heart of modern probability and the class of examples comprises the classical central limit theorem or the semicircular law in free probability. In this section, we shall derive a universality result for so-called homogeneous sums based on a sequence of independent centred Poisson random variables. For further background material concerning universality results for homogeneous sums we refer to the monograph [10] as well as to the original papers [11, 19].

We start by introducing the notion of a particularly well-behaved class of kernels.

Definition 4.1 (Index functions). Fix an integer $q \geq 1$. A function $h: \mathbb{N}^q \rightarrow \mathbb{R}$ is an *index function* of order q , if

- (a) h is symmetric meaning that $h(i_1, \dots, i_q) = h(i_{\pi(1)}, \dots, i_{\pi(q)})$ for all $(i_1, \dots, i_q) \in \mathbb{N}^q$ and all permutations $\pi \in \Pi_q$;
- (b) it vanishes on diagonals meaning that for $(i_1, \dots, i_q) \in \mathbb{N}^q$, $h(i_1, \dots, i_q) = 0$ whenever $i_k = i_\ell$ for some $k \neq \ell$.

Fix an integer $N \geq 1$. If g and h are two index functions of order q , we define their scalar product by

$$\langle g, h \rangle_{(N,q)} = \sum_{1 \leq i_1, \dots, i_q \leq N} g(i_1, \dots, i_q) h(i_1, \dots, i_q)$$

and write $\|h\|_{(N,q)} = \langle h, h \rangle_{(N,q)}^{1/2}$ for the corresponding norm. We frequently suppress the subscript (N, q) if it is clear from the context.

As in Section 3, we denote by $\{\mu_n : n \geq 1\}$ a sequence of σ -finite non-atomic measures on some Polish space $(\mathcal{Z}, \mathcal{Z})$.

Definition 4.2 (Tamed sequences). Fix an integer $q \geq 1$. A sequence $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ is *tamed* if there exists a sequence of integers $\{N_n : n \geq 1\}$ with $N_n \rightarrow \infty$, as $n \rightarrow \infty$, and an infinite measurable partition $\{B_i : i \geq 1\}$ of \mathcal{Z} verifying the following conditions:

- (a) there exists $\alpha \in (0, \infty)$ such that $\alpha < \mu_n(B_i) < \infty$ for every $i, n \geq 1$,
- (b) there is a sequence of index functions $\{h_n : n \geq 1\}$ of order q , such that f_n has the representation

$$f_n(z_1, \dots, z_q) = \sum_{1 \leq i_1, \dots, i_q \leq N_n} h_n(i_1, \dots, i_q) \prod_{k=1}^q \frac{\mathbf{1}_{B_{i_k}}(z_k)}{\sqrt{\mu_n(B_{i_k})}}. \quad (4.1)$$

Remark 4.3. (a) It follows from the definition that if a sequence $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ is tamed, we necessarily must have that $\mu_n(\mathcal{Z}) = \infty$ for every $n \geq 1$.

(b) If $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ is a tamed sequence with a representation as at (4.1), we have that $\|h_n\|_{(N_n,q)} = \|f_n\|_{L^2(\mu_n^q)} < \infty$.

(c) One easily verifies that tamed sequences automatically satisfy the technical assumptions (A).

Definition 4.4 (Homogeneous sums). Fix integers $N, q \geq 1$ and let $\mathbf{X} = \{X_i: i \geq 1\}$ be a sequence of random variables. Let h be an index function of order q . Then

$$Q_q(N, h, \mathbf{X}) = \sum_{1 \leq i_1, \dots, i_q \leq N} h(i_1, \dots, i_q) X_{i_1} \cdots X_{i_q}$$

is the *homogeneous sum* of h of order q based on the first N elements of \mathbf{X} .

If $\mathbf{X} = \{X_i: i \geq 1\}$ is a sequence of independent and centred random variables with unit variance, then

$$\mathbb{E}[Q_q(N, h, \mathbf{X})] = 0, \quad \mathbb{E}[Q_q(N, h, \mathbf{X})^2] = q! \|h\|_{(N, q)}^2.$$

In what follows, two particular classes of random variables play a special role. By $\mathbf{G} = \{G_i: i \geq 1\}$ we indicate a sequence of independent and identically distributed random variables, such that $G_i \sim \mathcal{N}(0, 1)$ for every $i \geq 1$. Moreover, we shall write $\mathbf{P} = \{P_i: i \geq 1\}$ for a sequence of independent random variables verifying

$$P_i \stackrel{d}{=} \frac{\text{Po}(\lambda_i) - \lambda_i}{\sqrt{\lambda_i}}, \quad i \geq 1,$$

where $\text{Po}(\lambda_i)$ indicates a Poisson random variable with mean λ_i , such that $\alpha = \inf\{\lambda_i: i \geq 1\} > 0$.

There is a close connection between homogeneous sums based on \mathbf{P} (or \mathbf{G}) and multiple stochastic integrals with respect to a centred Poisson measure $\hat{\eta}_n$ (or a Gaussian measure G_n) of tamed sequences. Namely, if $q \geq 1$ is a fixed integer and $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ is a tamed sequence with representation (4.1), then there is a sequence of centred Poisson measures $\{\hat{\eta}_n: n \geq 1\}$ (or a sequence of Gaussian measures $\{G_n: n \geq 1\}$) such that

$$I_q(f_n) = Q_q(N_n, h_n, \mathbf{P}), \quad I_q^{G_n}(f_n) = Q_q(N_n, h_n, \mathbf{G}). \quad (4.2)$$

Vice versa, given a sequence of index functions $\{h_n: n \geq 1\}$ of order $q \geq 1$ and a sequence of integers $\{N_n: n \geq 1\}$ diverging to infinity, as $n \rightarrow \infty$, such that $\|h_n\|_{(N_n, q)} < \infty$ for every $n \geq 1$, then there is a tamed sequence $\{f_n: n \geq 1\}$ with representation (4.1) and sequences of centred Poisson measures $\{\hat{\eta}_n: n \geq 1\}$ and Gaussian measures $\{G_n: n \geq 1\}$ such that (4.2) holds.

The following result is a version of [11, Theorem 1.8] and [11, Theorem 1.12]. Notice that there, the results are stated for integer-valued parameters $\nu \geq 1$, but they continue to hold for any $\nu > 0$.

Theorem 4.5 (Gamma universality of homogeneous sums on a fixed Gaussian chaos). *Fix $\nu > 0$, let $q \geq 2$ be even and $\{f_n: n \geq 1\} \subset L^2(\mu_n^q)$ be a tamed sequence with representation (4.1) that satisfies the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q^G(f_n)^2] = \lim_{n \rightarrow \infty} \mathbb{E}[Q_q(N_n, h_n, \mathbf{G})^2] = 2\nu.$$

Let $Y \sim \bar{\Gamma}_\nu$ be a centred Gamma random variable with parameter ν . Then the following five assertions are equivalent:

- (i) As $n \rightarrow \infty$, the sequence $\{Q_q(N_n, h_n, \mathbf{G}): n \geq 1\}$ converges in distribution to Y ;
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[Q_q(N_n, h_n, \mathbf{G})^4] - 12\mathbb{E}[Q_q(N_n, h_n, \mathbf{G})^3] = 12\nu^2 - 48\nu$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^r f_n\| = 0$ for every $r \in \{1, \dots, q-1\} \setminus \{q/2\}$, and

$$\lim_{n \rightarrow \infty} \|f_n \star_{q/2}^{q/2} f_n - c_q f_n\| = 0$$
 with $c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}$;

- (iv) for every sequence $\mathbf{X} = \{X_i: i \geq 1\}$ of independent centred random variables with unit variance which is such that $\sup_i \mathbb{E}|X_i|^{2+\varepsilon} < \infty$ for some $\varepsilon > 0$, the sequence $\{Q_q(N_n, h_n, \mathbf{X}): n \geq 1\}$ converges in distribution to Y , as $n \rightarrow \infty$;
- (v) for every sequence $\mathbf{X} = \{X_i: i \geq 1\}$ of i.i.d. centred random variables with unit variance, the sequence $\{Q_q(N_n, h_n, \mathbf{X}): n \geq 1\}$ converges in distribution to Y , as $n \rightarrow \infty$.

The following result answers the question whether Theorem 4.5 continues to hold if in (i) and (ii) the class \mathbf{G} is replaced by \mathbf{P} . Due to the discussion in Section 3.2, we cannot avoid additional assumptions in the Poisson case. In particular, we have to assume that either $q = 2$ or $q = 4$.

Theorem 4.6 (Gamma universality of homogeneous sums on a fixed Poisson chaos). *Fix $\nu > 0$ and let $q \geq 2$ be even and $\{f_n: n \geq 1\} \subset L^2(\mu_n^q)$ be a tamed sequence with representation (4.1) that satisfies the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = \lim_{n \rightarrow \infty} \mathbb{E}[Q_q(N_n, h_n, \mathbf{P})^2] = 2\nu. \quad (4.3)$$

Let $Y \sim \bar{\Gamma}_\nu$ be a random variable following a centred Gamma distribution with parameter ν . If one of the conditions

- (a) $q = 2$ and $\lim_{n \rightarrow \infty} \|f_n^2\| = 0$,
 (b) $q = 4$ and $f_n \leq 0$ for all $n \geq 1$

is satisfied, then the following five assertions are equivalent:

- (i) As $n \rightarrow \infty$, the sequence $\{Q_q(N_n, h_n, \mathbf{P}): n \geq 1\}$ converges in distribution to Y ;
- (ii) $\lim_{n \rightarrow \infty} \mathbb{E}[Q_q(N_n, h_n, \mathbf{P})^4] - 12\mathbb{E}[Q_q(N_n, h_n, \mathbf{P})^3] = 12\nu^2 - 48\nu$;
- (iii) $\lim_{n \rightarrow \infty} \|f_n \star_r^r f_n\| = 0$ for all $r \in \{1, \dots, q-1\} \setminus \{q/2\}$, and
 $\lim_{n \rightarrow \infty} \|f_n \widetilde{\star}_{q/2}^{q/2} f_n - c_q f_n\| = 0$ with $c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}$;
- (iv) for every sequence $\mathbf{X} = \{X_i: i \geq 1\}$ of independent centred random variables with unit variance which is such that $\sup_i \mathbb{E}|X_i|^{2+\varepsilon} < \infty$ for some $\varepsilon > 0$, the sequence $\{Q_q(N_n, h_n, \mathbf{X}): n \geq 1\}$ converges in distribution to Y , as $n \rightarrow \infty$;
- (v) for every sequence $\mathbf{X} = \{X_i: i \geq 1\}$ of i.i.d. centred random variables with unit variance, the sequence $\{Q_q(N_n, h_n, \mathbf{X}): n \geq 1\}$ converges in distribution to Y , as $n \rightarrow \infty$.

Proof. At first, we observe that due to Theorem 4.5, the assertions (iii), (iv) and (v) are equivalent. In [19, Subsection 4.2], it has been argued that

$$\sup_{i \geq 1} \mathbb{E}|P_i|^p < \infty \quad (4.4)$$

for all $p \geq 1$. This means that \mathbf{P} is a special instance of a sequence with the properties in assertion (iv) such that we obtain the implication (iv) \implies (i). Moreover, (4.4) implies together with the normalization condition (4.3) and [11, Lemma 4.2] that the sequence $\{Q_q(N_n, h_n, \mathbf{P})^4: n \geq 1\}$ is uniformly integrable such that we get the implication (i) \implies (ii). To prove (ii) \implies (iii), we apply Theorem 3.5. For this, one has to observe that assertion (iii) in Theorem 3.5 implies assertion (iii) in Theorem 4.6. \square

Remark 4.7. Theorem 4.6 shows that one can dispense with the assumption on the uniform integrability of the sequence $\{I_q(f_n)^4: n \geq 1\}$ in Theorem 3.5 whenever the sequence $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ is tamed.

Remark 4.8. Replacing in (b) the condition that $f_n \leq 0$ by $f_n \geq 0$, in (ii) the moment condition by $\lim_{n \rightarrow \infty} \mathbb{E}[Q_q(N_n, h_n, \mathbf{P})^4] + 12\mathbb{E}[Q_q(N_n, h_n, \mathbf{P})^3] = 12\nu^2 - 48\nu$ and in (iii) the condition on the middle-contraction by $\|f_n \tilde{\star}_{q/2}^{q/2} f_n + c_q f_n\| \rightarrow 0$, one arrives at a version of Theorem 4.6 with a centred reflected limiting random variable $Y \sim \widehat{\Gamma}_\nu$ in assertion (i), (iv) and (v).

4.2. U -statistics. Our second application is concerned with U -statistics. To introduce them, fix an integer $d \geq 1$ and let $\mathbf{Y} = \{Y_i : i \geq 1\}$ be a sequence of i.i.d. random vectors in \mathbb{R}^d , whose distribution has a density $p(\cdot)$ with respect to the Lebesgue measure on \mathbb{R}^d . Next, for any $n \geq 1$, let N_n be a Poisson random variable with mean n and define

$$\eta_n = \sum_{i=1}^{N_n} \delta_{Y_i}. \quad (4.5)$$

Clearly, η_n is a Poisson random measure on \mathbb{R}^d with control measure $\mu_n(dx) = np(x) dx$, implying that $\mu_n(\mathbb{R}^d) = n \rightarrow \infty$, as $n \rightarrow \infty$. Now, we put $\hat{\eta}_n = \eta_n - \mu_n$ and set $\mu = \mu_1$ for the sake of convenience. By a *Poisson U -statistic* of order $q \geq 2$ based on η_n we mean in this paper a random variable of the form

$$U_n = \sum_{1 \leq i_1 < \dots < i_q \leq N_n} h_n(Y_{i_1}, \dots, Y_{i_q}), \quad n \geq 1,$$

where the kernel $h_n: (\mathbb{R}^d)^q \rightarrow \mathbb{R}$ is an element of $L_s^1(\mu^q)$. On the other hand, a *classical U -statistic* is a random variable \hat{U}_n such that

$$\hat{U}_n = \sum_{1 \leq i_1 < \dots < i_q \leq n} h_n(Y_{i_1}, \dots, Y_{i_q}), \quad n \geq 1.$$

The difference between U_n and \hat{U}_n is that U_n involves a random number $\binom{N_n}{q}$ of summands, while the number of summands in the definition of \hat{U}_n is fixed (namely $\binom{n}{q}$). We say that a (Poisson or classical) U -statistic is *completely degenerate* if

$$\int_{\mathbb{R}^d} h_n(x, z_1, \dots, z_{q-1}) p(x) dx = 0$$

for μ^{q-1} -almost every $(z_1, \dots, z_{q-1}) \in (\mathbb{R}^d)^{q-1}$. In particular, this implies that $\mathbb{E}[U_n] = \mathbb{E}[\hat{U}_n] = 0$. Moreover, we suppose that U_n and \hat{U}_n are square-integrable.

We recall the following particular case of a celebrated theorem of de Jong, which provides a simple moment condition under which a central limit theorem for a sequence of completely degenerate U -statistics is guaranteed.

Theorem 4.9 (de Jong [2, 3]). *Let $q \geq 2$ and $\{h_n : n \geq 1\}$ be a sequence of non-zero elements of $L_s^4(\mu^q)$. Suppose that the U -statistics U_n and \hat{U}_n are completely degenerate and define $\sigma^2(n) = \text{Var}(U_n)$. Then the moment condition $\lim_{n \rightarrow \infty} \frac{\mathbb{E}[U_n^4]}{\sigma(n)^4} = 0$ implies that, as $n \rightarrow \infty$, the sequences $U_n/\sigma(n)$ and $\hat{U}_n/\sigma(n)$ converge in distribution to a standard Gaussian random variable.*

In our paper, we are interested in the Gamma approximation of Poisson and classical U -statistics. The next result generalizes Theorem 2.13 (B) in [17], where the authors had to restrict to the case $q = 2$. Here, we add a corresponding limit theorem in case that $q = 4$

under an additional sign condition. It can be seen as a non-central version of de Jong's theorem, Theorem 4.9. We shall see that in the non-central case a similar result is true under a suitable condition involving only the third and the fourth moment.

Theorem 4.10. *Suppose that $q \in \{2, 4\}$. For each $n \geq 1$ let $h_n \in L_s^4(\mu^q)$ be a function such that*

$$\sup_{n \geq 1} \frac{\int h_n^4 d\mu_n^q}{(\int h_n^2 d\mu_n^q)^2} < \infty$$

and suppose that the U -statistics U_n and \hat{U}_n are completely degenerate. Further assume that there exists $\nu > 0$ such that $\lim_{n \rightarrow \infty} \mathbb{E}[U_n^2] = 2\nu$ and that

- (a) $\lim_{n \rightarrow \infty} \|h_n^2\| = 0$ if $q = 2$,
- (b) $f_n \leq 0$ for all $n \geq 1$ if $q = 4$.

Then the moment condition $\lim_{n \rightarrow \infty} \mathbb{E}[U_n^4] - 12\mathbb{E}[U_n^3] = 12\nu^2 - 48\nu$ implies that both random variables U_n and \hat{U}_n converge in distribution to $Y \sim \bar{\Gamma}_\nu$, as $n \rightarrow \infty$.

Proof. Using the fact that the Poisson U -statistics U_n is an element of the sum of the first q Poisson chaoses with respect to $\hat{\eta}_n$ as introduced after (4.5) (see [20, Theorem 3.6]), as well as the fact that U_n is completely degenerate, one obtains that $U_n = I_q(h_n)$ for every $n \geq 1$. The result for the Poisson U -statistics U_n then follows immediately from Theorem 3.5. Moreover, it is known from [4] that $\mathbb{E}[(U_n - \hat{U}_n)^2] = O(n^{-1/2})$, as $n \rightarrow \infty$. This yields the result also for \hat{U}_n . \square

Remark 4.11. Using Theorem 2.6 in [17] or its generalization Proposition 5.1 below, one can add a rate of convergence (for a certain smooth probability distance) between U_n or \hat{U}_n and the limiting random variable Y . However, we do not pursue such quantitative results in this paper.

Remark 4.12. In assumption (b) of Theorem 4.10 one can replace the sign condition $f_n \leq 0$ by $f_n \geq 0$ and at the same time the moment condition $\mathbb{E}[U_n^4] - 12\mathbb{E}[U_n^3] \rightarrow 12\nu^2 - 48\nu$ by $\mathbb{E}[U_n^4] + 12\mathbb{E}[U_n^3] \rightarrow 12\nu^2 - 48\nu$. In this case, the limiting random variable Y has a centred reflected Gamma distribution $\hat{\Gamma}_\nu$ with parameter $\nu > 0$.

5. PROOF OF THEOREM 3.5

5.1. Strategy of the proof. Before entering the details of the proof of Theorem 3.5, let us briefly summarize the overall strategy.

First of all, the implication (i) \implies (ii) of Theorem 3.5 is a direct consequence of the uniform integrability of the sequence $\{I_q(f_n)^4 : n \geq 1\}$. Next, the implication (iii) \implies (i) will follow from a generalization of the main result of [17], which has been derived by the Malliavin-Stein method. It delivers a criterion in terms of contraction norms, which ensures centred Gamma convergence on a fixed Poisson chaos of even order and is presented as Proposition 5.1 below. The main part of proof of Theorem 3.5 consists in showing that (ii) implies (iii). It is based on the technical Lemmas 5.2 and 5.4, which establish new inequalities for norms of contraction kernels, that are also of independent interest. Next, in Lemma 5.6 we derive an asymptotic lower bound for the moment expression $\mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3]$ in terms of contraction norms. Finally, Lemma 5.7 shows under the conditions of Theorem 3.5 that if the lower bound for $\mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3]$ converges to the 'correct' quantity, the contraction conditions in (iii) are satisfied. Lemma 5.9 proves that this lower bound actually converges.

We emphasize that we state all intermediate steps of the proof of Theorem 3.5 as general as possible in order to highlight in which step the restrictive condition that $q = 2$ or $q = 4$ and the sign condition on the kernels arise.

5.2. Preparatory steps. We start our investigations with a generalization of Theorem 2.6 in [17]. The main difference between that result and Proposition 5.1 is that for technical reasons it has been assumed in [17] that μ_n is a finite measure for each $n \geq 1$ such that $\mu_n(\mathcal{Z}) \rightarrow \infty$, as $n \rightarrow \infty$. Our next result shows that one can dispense with this assumption.

Proposition 5.1. *Fix $\nu > 0$ and an even integer $q \geq 2$. Let the sequence $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ satisfy the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = 2\nu.$$

Then, if

$$\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0 \text{ for all } r \in \{1, \dots, q\} \text{ and } \ell \in \{1, \dots, r \wedge (q-1)\}, (r, \ell) \neq (q/2, q/2),$$

$$\lim_{n \rightarrow \infty} \|f_n^2\| = 0,$$

$$\lim_{n \rightarrow \infty} \|f_n \tilde{\star}_{q/2}^{q/2} f_n - c_q f_n\| = 0 \text{ with } c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2},$$

the sequence $\{I_q(f_n): n \geq 1\}$ converges in distribution to $Y \sim \bar{\Gamma}_\nu$, as $n \rightarrow \infty$.

Proof. In principle, one can follow the proof of [17, Theorem 2.6]. The only part where the assumption about the finiteness of the measures μ_n enters is [17, Proposition 2.3]. To circumvent this problem, one uses the modified integration-by-parts formula [21, Lemma 2.3] and concludes as in the proof of Theorem 4.1 of [5]. Since the computations are quite straight forward, we omit the details. \square

We now present two estimates of the norm of a symmetrized contraction kernel in terms of non-symmetrized contraction norms. In particular, our first lemma generalizes [16, Identity (11.6.30)]. We recall for $f \in L_s^2(\mu^q)$, $q \geq 1$, that $\|f \tilde{\star}_q^q f\|^2 = \|f \star_q^q f\|^2 = \|f\|^4$ and $\|f \star_0^0 f\|^2 = \|f\|^4$.

Lemma 5.2. *Let $q \geq 1$ be an integer and $f \in L_s^2(\mu^q)$ be a kernel satisfying the technical assumptions (A). Then*

$$\|f \tilde{\star}_0^0 f\|^2 = \frac{(q!)^2}{(2q)!} \left(2\|f\|^4 + \sum_{p=1}^{q-1} \binom{q}{p}^2 \|f \star_p^p f\|^2 \right). \quad (5.1)$$

Furthermore, for any $r \in \{1, \dots, q-1\}$ one has the inequality

$$\|f \tilde{\star}_r^r f\|^2 \leq \frac{((q-r)!)^2}{(2(q-r))!} \left(2\|f \star_r^r f\|^2 + \sum_{p=1}^{q-r-1} \binom{q-r}{p}^2 \|f \star_p^p f\|^2 \right). \quad (5.2)$$

If $q \geq 2$ is an even integer, Equation (5.2) yields that

$$\|f \tilde{\star}_{q/2}^{q/2} f\|^2 \leq \frac{((q/2)!)^2}{q!} \left(2\|f \star_{q/2}^{q/2} f\|^2 + \sum_{p=1}^{q/2-1} \binom{q/2}{p}^2 \|f \star_p^p f\|^2 \right). \quad (5.3)$$

This inequality will turn out to be crucial in what follows.

Before entering the proof of Lemma 5.2, we introduce some notation. Recall that for an integer $p \geq 1$, we denote the group of $p!$ permutations of the set $\{1, \dots, p\}$ by Π_p . For a kernel $g \in L^2(\mu^p)$ and a permutation $\pi \in \Pi_p$, we use the shorthand $g(\pi)$ for the mapping $\mathcal{Z}^p \ni (z_1, \dots, z_p) \mapsto g(\pi)(z_1, \dots, z_p) = g(z_{\pi(1)}, \dots, z_{\pi(p)})$. We can immediately see that $\|g\| = \|g(\pi)\|$ for all $\pi \in \Pi_p$ such that automatically $g(\pi) \in L^2(\mu^p)$. In the following, we use the convention that $\pi_0 \in \Pi_p$ is the identity map, meaning that $g(\pi_0) = g$.

For any integer $M \geq 1$, any two permutations $\pi, \sigma \in \Pi_{2M}$ and any $p \in \{0, \dots, M\}$ we shall use the notation

$$\pi \sim_p \sigma$$

if and only if

$$|\{\pi(1), \dots, \pi(M)\} \cap \{\sigma(1), \dots, \sigma(M)\}| = p,$$

where $|\cdot|$ stands for the cardinality of the argument set. If $\pi \sim_p \sigma$, then clearly $|\{\pi(M+1), \dots, \pi(2M)\} \cap \{\sigma(M+1), \dots, \sigma(2M)\}| = p$. In the proof of [16, Proposition 11.2.2], there is an explanation that, given a permutation $\pi \in \Pi_{2M}$ and an integer $p \in \{0, \dots, M\}$, there are exactly $(M!)^2 \binom{M}{p}^2$ permutations $\sigma \in \Pi_{2M}$ such that $\pi \sim_p \sigma$.

Proof of Lemma 5.2. Let $q \geq 1$ be an integer and $f \in L_s^2(\mu^q)$ be a kernel satisfying the technical assumptions (A). Fix $r \in \{0, 1, \dots, q-1\}$. We have that

$$\begin{aligned} \|f \star_r^r f\|^2 &= \langle f \star_r^r f, f \star_r^r f \rangle = \frac{1}{(2q-2r)!} \sum_{\pi \in \Pi_{2q-2r}} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \\ &= \frac{1}{(2q-2r)!} \sum_{p=0}^{q-r} \sum_{\pi \sim_p \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle. \end{aligned} \quad (5.4)$$

To prove (5.1), let $r = 0$ and $\pi \sim_0 \pi_0$ or $\pi \sim_q \pi_0$. Then we get

$$\begin{aligned} &\langle f \star_0^0 f, f \star_0^0 f(\pi) \rangle \\ &= \int_{\mathcal{Z}^{2q}} f(z_1, \dots, z_q) f(z_{q+1}, \dots, z_{2q}) f(z_{\pi(1)}, \dots, z_{\pi(q)}) f(z_{\pi(q+1)}, \dots, z_{\pi(2q)}) \mu^{2q}(\mathrm{d}(z_1 \dots z_{2q})) \\ &= \left(\int_{\mathcal{Z}^q} f(w_1, \dots, w_q)^2 \mu^q(\mathrm{d}(w_1 \dots w_q)) \right)^2 \\ &= \|f\|^4. \end{aligned}$$

Now, let $\pi \sim_p \pi_0$ with $p \in \{1, \dots, q-1\}$. Then

$$\begin{aligned} &\langle f \star_0^0 f, f \star_0^0 f(\pi) \rangle \\ &= \int_{\mathcal{Z}^{2q}} f(z_1, \dots, z_q) f(z_{q+1}, \dots, z_{2q}) f(z_{\pi(1)}, \dots, z_{\pi(q)}) f(z_{\pi(q+1)}, \dots, z_{\pi(2q)}) \mu^{2q}(\mathrm{d}(z_1 \dots z_{2q})) \\ &= \int_{\mathcal{Z}^{2q-2p} \times \mathcal{Z}^p \times \mathcal{Z}^p} f(z_1, \dots, z_q) f(z_{\pi(1)}, \dots, z_{\pi(q)}) \\ &\quad \times f(z_{q+1}, \dots, z_{2q}) f(z_{\pi(q+1)}, \dots, z_{\pi(2q)}) \mu^{2q}(\mathrm{d}(z_1 \dots z_{2q})) \\ &\stackrel{(*)}{=} \int_{\mathcal{Z}^{2q-2p}} f \star_p^p f(w_1, \dots, w_{2q-2p}) \times f \star_p^p f(w_1, \dots, w_{2q-2p}) \mu^{2q-2p}(\mathrm{d}(w_1 \dots w_{2q-2p})) \\ &= \|f \star_p^p f\|^2. \end{aligned}$$

We note that the assumption that f is symmetric is essential to get the identity highlighted by (\star) . In view of (5.4), we obtain

$$\begin{aligned} & \|f \star_0^0 f\|^2 \\ &= \frac{1}{(2q)!} \left(\sum_{\pi \sim_0 \pi_0} \langle f \star_0^0 f, f \star_0^0 f(\pi) \rangle + \sum_{\pi \sim_q \pi_0} \langle f \star_0^0 f, f \star_0^0 f(\pi) \rangle + \sum_{p=1}^{q-1} \sum_{\pi \sim_p \pi_0} \langle f \star_0^0 f, f \star_0^0 f(\pi) \rangle \right) \\ &= \frac{1}{(2q)!} \left(2(q!)^2 \|f\|^4 + \sum_{p=1}^{q-1} (q!)^2 \binom{q}{p}^2 \|f \star_p^p f\|^2 \right), \end{aligned}$$

such that (5.1) follows. Now, let $r \in \{1, \dots, q-1\}$ and observe that for $\pi \sim_{q-r} \pi_0$ one has that

$$\begin{aligned} & \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \\ &= \int_{\mathcal{Z}^{2q-2r}} \left(\int_{\mathcal{Z}^r} f(x_1, \dots, x_r, z_1, \dots, z_{q-r}) f(x_1, \dots, x_r, z_{q-r+1}, \dots, z_{2q-2r}) \mu^r(d(x_1 \dots x_r)) \right) \\ & \quad \times \left(\int_{\mathcal{Z}^r} f(y_1, \dots, y_r, z_{\pi(1)}, \dots, z_{\pi(q-r)}) f(y_1, \dots, y_r, z_{\pi(q-r+1)}, \dots, z_{\pi(2q-2r)}) \mu^r(d(y_1 \dots y_r)) \right) \\ & \quad \mu^{2q-2r}(d(z_1 \dots z_{2q-2r})) \\ &= \int_{\mathcal{Z}^{2q}} f(x_1, \dots, x_r, z_1, \dots, z_{q-r}) f(y_1, \dots, y_r, z_{\pi(1)}, \dots, z_{\pi(q-r)}) \\ & \quad \times f(x_1, \dots, x_r, z_{q-r+1}, \dots, z_{2q-2r}) f(y_1, \dots, y_r, z_{\pi(q-r+1)}, \dots, z_{\pi(2q-2r)}) \\ & \quad \mu^{2q}(d(x_1 \dots x_r, y_1 \dots y_r, z_1 \dots z_{2q-2r})) \\ &= \int_{\mathcal{Z}^{2r} \times \mathcal{Z}^{q-r} \times \mathcal{Z}^{q-r}} f(x_1, \dots, x_r, z_1, \dots, z_{q-r}) f(y_1, \dots, y_r, z_{\pi(1)}, \dots, z_{\pi(q-r)}) \\ & \quad \times f(x_1, \dots, x_r, z_{q-r+1}, \dots, z_{2q-2r}) f(y_1, \dots, y_r, z_{\pi(q-r+1)}, \dots, z_{\pi(2q-2r)}) \\ & \quad \mu^{2q}(d(x_1 \dots x_r, y_1 \dots y_r, z_1 \dots z_{2q-2r})) \\ &= \int_{\mathcal{Z}^{2r}} \left(f \star_{q-r}^{q-r} f(x_1, \dots, x_r, y_1, \dots, y_r) \right)^2 \mu^{2r}(d(x_1 \dots x_r, y_1 \dots y_r)) \\ &= \|f \star_{q-r}^{q-r} f\|^2 \\ &= \|f \star_r^r f\|^2. \end{aligned}$$

Similarly, we obtain for the case that $\pi \sim_0 \pi_0$,

$$\langle f \star_r^r f, f \star_r^r f(\pi) \rangle = \|f \star_r^r f\|^2.$$

Now, let $\pi \sim_p \pi_0$ with $p \in \{1, \dots, q-r-1\}$. Then, there is a permutation $\sigma \in \Pi_{2q-2p}$ such that

$$\begin{aligned}
& \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \tag{5.5} \\
&= \int_{\mathcal{Z}^{2q}} f(x_1, \dots, x_r, z_1, \dots, z_{q-r}) f(y_1, \dots, y_r, z_{\pi(1)}, \dots, z_{\pi(q-r)}) \\
&\quad \times f(x_1, \dots, x_r, z_{q-r+1}, \dots, z_{2q-2r}) f(y_1, \dots, y_r, z_{\pi(q-r+1)}, \dots, z_{\pi(2q-2r)}) \\
&\quad \mu^{2q}(\mathrm{d}(x_1 \dots x_r, y_1 \dots y_r, z_1 \dots z_{2q-2r})) \\
&= \int_{\mathcal{Z}^{2q-2p} \times \mathcal{Z}^p \times \mathcal{Z}^p} f(x_1, \dots, x_r, z_1, \dots, z_{q-r}) f(y_1, \dots, y_r, z_{\pi(1)}, \dots, z_{\pi(q-r)}) \\
&\quad \times f(x_1, \dots, x_r, z_{q-r+1}, \dots, z_{2q-2r}) f(y_1, \dots, y_r, z_{\pi(q-r+1)}, \dots, z_{\pi(2q-2r)}) \\
&\quad \mu^{2q}(\mathrm{d}(x_1 \dots x_r, y_1 \dots y_r, z_1 \dots z_{2q-2r})) \\
&= \int_{\mathcal{Z}^{2q-2p}} f \star_p^p f(w_1, \dots, w_{2q-2p}) \times f \star_p^p f(w_{\sigma(1)}, \dots, w_{\sigma(2q-2p)}) \mu^{2q-2p}(\mathrm{d}(w_1 \dots w_{2q-2p})) \\
&= \langle f \star_p^p f, f \star_p^p f(\sigma) \rangle \\
&\leq \|f \star_p^p f\| \|f \star_p^p f(\sigma)\| \\
&= \|f \star_p^p f\|^2.
\end{aligned}$$

Note that contrary to the case $r = 0$, σ shows up because of the appearance of the variables $x_1, \dots, x_r, y_1, \dots, y_r$. Therefore, we need to apply the Cauchy-Schwarz inequality once, which is the very reason for the inequality in (5.2). At this stage, (5.2) follows by (5.4) and

$$\begin{aligned}
\|f \widetilde{\star}_r^r f\|^2 &= \frac{1}{(2q-2r)!} \left(\sum_{\pi \sim_0 \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle + \sum_{\pi \sim_{q-r} \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \right. \\
&\quad \left. + \sum_{p=1}^{q-r-1} \sum_{\pi \sim_p \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \right) \\
&\leq \frac{1}{(2q-2r)!} \left(2((q-r)!)^2 \|f \star_r^r f\|^2 + \sum_{p=1}^{q-r-1} ((q-r)!)^2 \binom{q-r}{p}^2 \|f \star_p^p f\|^2 \right).
\end{aligned}$$

This completes the proof. \square

Remark 5.3. A combinatorial argument shows that the permutation $\sigma \in \Pi_{2q-2p}$ appearing in (5.5) cannot be such that $f \star_p^p f(\sigma) = f \star_p^p f$ (in particular, σ cannot be the identity). Hence, we cannot omit applying Cauchy-Schwarz in this case.

In Lemma 5.2 no condition on the sign of f was necessary. However, if we assume that f has constant sign, we are able to deduce a ‘reverse’ counterpart of (5.2).

Lemma 5.4. *Let $q \geq 1$ be an integer and $f \in L_s^2(\mu^q)$ a kernel satisfying the technical assumptions (A). If $f \leq 0$ or $f \geq 0$, then, for any $r \in \{0, 1, \dots, q-1\}$, one has that*

$$\|f \widetilde{\star}_r^r f\|^2 \geq \frac{2((q-r)!)^2}{(2q-2r)!} \|f \star_r^r f\|^2. \tag{5.6}$$

Proof. The left-hand side of (5.6) satisfies identity (5.4). Using the fact that f has constant sign, the right-hand side of (5.4) becomes smaller if we sum only over a subset of Π_{2q-2r} , namely over all $\pi \in \Pi_{2q-2r}$ such that $\pi \sim_0 \pi_0$ or $\pi \sim_{q-r} \pi_0$. Hence, we end up with

$$\begin{aligned} \|f \widetilde{\star}_r^r f\|^2 &\geq \frac{1}{(2q-2r)!} \left(\sum_{\pi \sim_0 \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle + \sum_{\pi \sim_{q-r} \pi_0} \langle f \star_r^r f, f \star_r^r f(\pi) \rangle \right) \\ &= \frac{2((q-r)!)^2}{(2(q-r))!} \|f \star_r^r f\|^2, \end{aligned}$$

which completes the proof. \square

Remark 5.5. In view of Remark 5.3, inequality (5.6) is optimal under the conditions of Lemma 5.4.

5.3. Proof of the implication (ii) \implies (iii). Let us introduce some notation. We shall write $a_n \asymp b_n$ for two real-valued sequences $\{a_n: n \geq 1\}$, $\{b_n: n \geq 1\}$, whenever $\lim_{n \rightarrow \infty} a_n - b_n = 0$. Be aware that this does not necessarily imply that one of the individual sequences converges, but of course ensures the convergence of both sequences whenever one of them converges. The next lemma establishes an asymptotic lower bound for the linear combination of the fourth and third moment $\mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3]$ of a sequence of Poisson integrals of even order $q \geq 2$ where $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$. It is one of the main ingredients to show the implication (ii) \implies (iii) in Theorem 3.5. Note that this bound holds for general even $q \geq 2$. Moreover, at this point we do not need an assumption on the sign of the kernels.

Lemma 5.6. *Let $\nu > 0$ and $q \geq 2$ be an even integer. Let $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence of kernels such that the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = 2\nu$$

are satisfied. Then one has that

$$\mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3] \asymp 12\nu^2 - 48\nu + A(I_q(f_n)) + R(I_q(f_n)), \quad (5.7)$$

where the terms on the right-hand side of (5.7) satisfy $A(I_q(f_n)) \geq A'(I_q(f_n))$ with

$$\begin{aligned} A'(I_q(f_n)) &= \sum_{p=1}^{q/2-1} \frac{(q!)^4}{(p!)^2} \left(\frac{2}{(q-p)!^2} - \frac{1}{2((q/2)!(q/2-p)!)^2} \right) \|f_n \star_p^p f_n\|^2 \\ &\quad + \sum_{p=1, p \neq q}^{2q-1} p! \|G_p^q f_n\|^2 + q! \sum_{p=q/2+1}^q (p!)^2 \binom{p}{q}^4 \binom{p}{q-p}^2 \|f_n \widetilde{\star}_p^{q-p} f_n\|^2 \\ &\quad + 24q! \|c_q^{-1} f_n \widetilde{\star}_{q/2}^{q/2} f_n - f_n\|^2 \end{aligned} \quad (5.8)$$

with $c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}$, and

$$\begin{aligned} R(I_q(f_n)) &= q! \sum_{\substack{r,p=q/2 \\ r \neq p}}^q r! p! \binom{q}{r}^2 \binom{q}{p}^2 \binom{r}{q-r} \binom{p}{q-p} \langle f_n \tilde{\star}_r^{q-r} f_n, f_n \tilde{\star}_p^{q-p} f_n \rangle \\ &\quad - 12q! \sum_{p=q/2+1}^q p! \binom{q}{p}^2 \binom{p}{q-p} \langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle. \end{aligned} \quad (5.9)$$

Proof of Lemma 5.6. In view of Lemma 2.2 and since q is even, one has that

$$\begin{aligned} &\mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3] \\ &= \sum_{p=0}^{2q} p! \|G_p^q f_n\|^2 - 12q! \sum_{p=q/2}^q p! \binom{q}{p}^2 \binom{p}{q-p} \langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle \\ &= (q!)^2 \|f_n\|^4 + (2q)! \|f_n \tilde{\star}_0^0 f_n\|^2 + \sum_{p=1}^{2q-1} p! \|G_p^q f_n\|^2 \\ &\quad - 12q! \sum_{p=q/2}^q p! \binom{q}{p}^2 \binom{p}{q-p} \langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle \\ &= 3(q!)^2 \|f_n\|^4 + \sum_{p=1}^{q-1} \frac{(q!)^4}{(p!(q-p)!)^2} \|f_n \star_p^p f_n\|^2 + \sum_{p=1}^{2q-1} p! \|G_p^q f_n\|^2 \\ &\quad - 12q! \sum_{p=q/2}^q p! \binom{q}{p}^2 \binom{p}{q-p} \langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle \\ &= 3(q!)^2 \|f_n\|^4 + T_1(I_q(f_n)) + T_2(I_q(f_n)) + T_3(I_q(f_n)), \end{aligned}$$

where the third equality stems from (5.1). The terms T_1, T_2, T_3 read as follows:

$$\begin{aligned} T_1(I_q(f_n)) &= \sum_{\substack{p=1 \\ p \neq q/2}}^{q-1} \frac{(q!)^4}{(p!(q-p)!)^2} \|f_n \star_p^p f_n\|^2 + \sum_{p=1, p \neq q}^{2q-1} p! \|G_p^q f_n\|^2, \\ T_2(I_q(f_n)) &= \frac{(q!)^4}{(q/2)!^4} \|f_n \star_{q/2}^{q/2} f_n\|^2 + q! \|G_{q/2}^q f_n\|^2 - 12q! (q/2)! \binom{q}{q/2}^2 \langle f_n \tilde{\star}_{q/2}^{q/2} f_n, f_n \rangle, \\ T_3(I_q(f_n)) &= -12q! \sum_{p=q/2+1}^q p! \binom{q}{p}^2 \binom{p}{q-p} \langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle. \end{aligned}$$

We use (5.3) to see that

$$\frac{(q!)^4}{(q/2)!^4} \|f_n \star_{q/2}^{q/2} f_n\|^2 \geq \frac{(q!)^5}{2(q/2)!^6} \|f_n \tilde{\star}_{q/2}^{q/2} f_n\|^2 - \frac{1}{2} \sum_{p=1}^{q/2-1} \frac{(q!)^4}{((q/2)! p! (q/2-p)!)^2} \|f_n \star_p^p f_n\|^2.$$

Using the definition of $G_q^q f_n$ given at (2.5), we have the estimate

$$\begin{aligned}
T_2(I_q(f_n)) &\geq q! \left(\left\| \sum_{r=q/2}^q r! \binom{q}{r}^2 \binom{r}{q-r} f_n \tilde{\star}_r^{q-r} f_n \right\|^2 + \frac{1}{2} \frac{(q!)^4}{(q/2)!^6} \|f_n \tilde{\star}_{q/2}^{q/2} f_n\|^2 \right. \\
&\quad \left. - 12(q/2)! \binom{q}{q/2}^2 \langle f_n \tilde{\star}_{q/2}^{q/2} f_n, f_n \rangle \right) - \frac{1}{2} \sum_{p=1}^{q/2-1} \frac{(q!)^4}{((q/2)!p!(q/2-p)!)^2} \|f_n \star_p^p f_n\|^2 \\
&= q! \left(\frac{3}{2} \frac{(q!)^4}{(q/2)!^6} \|f_n \tilde{\star}_{q/2}^{q/2} f_n\|^2 - 12 \frac{(q!)^2}{(q/2)!^3} \langle f_n \tilde{\star}_{q/2}^{q/2} f_n, f_n \rangle \right) \\
&\quad + q! \sum_{r=q/2+1}^q (r!)^2 \binom{q}{r}^4 \binom{r}{q-r}^2 \|f_n \tilde{\star}_r^{q-r} f_n\|^2 \\
&\quad + q! \sum_{\substack{r,p=q/2 \\ r \neq p}}^q r!p! \binom{q}{r}^2 \binom{q}{p}^2 \binom{r}{q-r} \binom{p}{q-p} \langle f_n \tilde{\star}_r^{q-r} f_n, f_n \tilde{\star}_p^{q-p} f_n \rangle \\
&\quad - \frac{1}{2} \sum_{p=1}^{q/2-1} \frac{(q!)^4}{((q/2)!p!(q/2-p)!)^2} \|f_n \star_p^p f_n\|^2.
\end{aligned}$$

Using the relation $\|f_n \star_p^p f_n\| = \|f_n \star_{q-p}^{q-p} f_n\|$, valid for all $p \in \{1, \dots, q-1\}$, we obtain

$$\begin{aligned}
&\sum_{\substack{p=1 \\ p \neq q/2}}^{q-1} \frac{(q!)^4}{(p!(q-p)!)^2} \|f_n \star_p^p f_n\|^2 - \frac{1}{2} \sum_{p=1}^{q/2-1} \frac{(q!)^4}{((q/2)!p!(q/2-p)!)^2} \|f_n \star_p^p f_n\|^2 \\
&= \sum_{p=1}^{q/2-1} \frac{(q!)^4}{(p!)^2} \left(\frac{2}{(q-p)!^2} - \frac{1}{2((q/2)!p!(q/2-p)!)^2} \right) \|f_n \star_p^p f_n\|^2.
\end{aligned}$$

The proof is concluded by observing that

$$\begin{aligned}
&q! \left(\frac{3}{2} \frac{(q!)^4}{((q/2)!^6} \|f_n \tilde{\star}_{q/2}^{q/2} f_n\|^2 - 12 \frac{(q!)^2}{((q/2)!^3} \langle f_n \tilde{\star}_{q/2}^{q/2} f_n, f_n \rangle \right) \\
&= \frac{3}{2} q! \left(\frac{(q!)^4}{((q/2)!^6} \|f_n \tilde{\star}_{q/2}^{q/2} f_n\|^2 - 2 \times 4 \frac{(q!)^2}{((q/2)!^3} \langle f_n \tilde{\star}_{q/2}^{q/2} f_n, f_n \rangle + 16 \|f_n\|^2 \right) - 24q! \|f_n\|^2 \\
&= 24q! \|c_q^{-1} f_n \tilde{\star}_{q/2}^{q/2} f_n - f_n\|^2 - 24q! \|f_n\|^2
\end{aligned}$$

and by recalling condition (a), which implies that $3(q!)^2 \|f_n\|^4 - 24q! \|f_n\|^2 \rightarrow 12\nu^2 - 48\nu$. \square

While all previous results did not use the assumptions on the order of the integral and the sign of the kernels, in the next lemma we need that $q \in \{2, 4\}$ and that the kernels have constant sign.

Lemma 5.7. *Let $\nu > 0$ and $q \in \{2, 4\}$. Let $\{f_n : n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence of kernels such that the technical assumptions (A) and the normalization condition $\lim_{n \rightarrow \infty} q! \|f_n\|^2 = 2\nu$ are satisfied. Assume that for each $n \geq 1$ either $f_n \leq 0$ or $f_n \geq 0$. Then the following two assertions concerning the term $A'(I_q(f_n))$ defined at (5.8) are true:*

- (1) $A'(I_q(f_n)) \geq 0$ for all $n \geq 1$;
(2) If $A'(I_q(f_n)) \rightarrow 0$, as $n \rightarrow \infty$, then

$$\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0 \quad (5.10)$$

for all $r \in \{1, \dots, q\}$ and $\ell \in \{1, \dots, r \wedge (q-1)\}$ such that $(r, \ell) \neq (q/2, q/2)$,

$$\lim_{n \rightarrow \infty} \|f_n^2\| = 0, \quad (5.11)$$

$$\lim_{n \rightarrow \infty} \|f_n \star_{q/2}^{\sim q/2} f_n - c_q f_n\| = 0 \quad \text{with } c_q = \frac{4}{(q/2)! \binom{q}{q/2}}. \quad (5.12)$$

Proof. We start by showing the first assertion of the lemma. The only term that might be negative on right-hand side of (5.8) is the first sum. For the case $q = 2$, this does not play any role, because then the sum vanishes. Hence, $A'(I_q(f_n))$ is a positive linear combination of non-negative terms.

Now, let $q \geq 4$ be even. Using the fact that f_n has constant sign, $\|f_n \star_p^p f_n\| = \|f_n \star_{q-p}^{q-p} f_n\|$ for all $p \in \{1, \dots, q-1\}$ as well as Lemma 5.4, we obtain the estimate

$$\begin{aligned} \sum_{p=1, p \neq q}^{2q-1} p! \|G_p^q f_n\|^2 &\geq \sum_{p=1, p \neq q}^{2q-1} p! \sum_{r=0}^q \sum_{\ell=0}^r \mathbf{1}(2q-r-\ell=p) r!^2 \binom{q}{r}^4 \binom{r}{\ell}^2 \|f_n \star_r^\ell f_n\|^2 \\ &\geq \sum_{\substack{p=1, p \neq q, \\ p \text{ even}}}^{2q-1} p! ((q-p/2)!)^2 \binom{q}{q-p/2}^4 \|f_n \star_{q-p/2}^{\sim q-p/2} f_n\|^2 \\ &= \sum_{p=1, p \neq q/2}^{q-1} (2p)! ((q-p)!)^2 \binom{q}{q-p}^4 \|f_n \star_{q-p}^{q-p} f_n\|^2 \\ &= \sum_{p=1, p \neq q/2}^{q-1} (2(q-p))! (p!)^2 \binom{q}{p}^4 \|f_n \star_p^p f_n\|^2 \\ &\geq \sum_{p=1, p \neq q/2}^{q-1} 2((q-p)!)^2 (p!)^2 \binom{q}{p}^4 \|f_n \star_p^p f_n\|^2 \\ &= \sum_{p=1}^{q/2-1} \frac{4(q!)^4}{((q-p)!)^2 (p!)^2} \|f_n \star_p^p f_n\|^2. \end{aligned}$$

Hence, we end up with

$$\begin{aligned} &\sum_{p=1}^{q/2-1} \frac{(q!)^4}{(p!)^2} \left(\frac{2}{(q-p)!^2} - \frac{1}{2((q/2)!(q/2-p)!)^2} \right) \|f_n \star_p^p f_n\|^2 + \sum_{p=1, p \neq q}^{2q-1} p! \|G_p^q f_n\|^2 \\ &\geq \sum_{p=1}^{q/2-1} \frac{(q!)^4}{(p!)^2} \left(\frac{6}{(q-p)!^2} - \frac{1}{2((q/2)!(q/2-p)!)^2} \right) \|f_n \star_p^p f_n\|^2. \end{aligned} \quad (5.13)$$

For $q = 4, p = 1$ we have that

$$\frac{6}{(q-p)!^2} - \frac{1}{2((q/2)!(q/2-p)!)^2} = \frac{1}{24} > 0.$$

So, for $q = 4$ (and $q = 2$), the term $A'(I_q(f_n))$ is bounded from below by a linear combination with positive coefficients of the norms of the contraction kernels appearing in (5.10), (5.11) and (5.12) (while for all even $q \geq 6$ this cannot be guaranteed any more). This proves both statements of the lemma. \square

Remark 5.8. As anticipated, for all even $q \geq 6$ there are combinatorial coefficients in (5.13) which are negative, implying that our proof cannot be generalized to Poisson integrals of arbitrary order. The reason is that one would need a sharper version of Lemma 5.4, which is in general not available as discussed in Remark 5.5. As a consequence, we have to leave it as an open problem to establish a four moments theorem for the Gamma approximation for Poisson integrals of order $q \geq 6$ by different methods.

It remains to check whether the conditions of Theorem 3.5 are sufficient to imply that $A'(I_q(f_n)) \rightarrow 0$. The following lemma shows that this is indeed the case.

Lemma 5.9. *Let $\nu > 0$ and $q \in \{2, 4\}$. Let $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^q)$ be a sequence of kernels satisfying the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} q! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^2] = 2\nu.$$

Let the sequence $\{I_q(f_n)^4: n \geq 1\}$ be uniformly integrable. If one of the conditions

- (a) $q = 2$ and $\lim_{n \rightarrow \infty} \|f_n^2\| = 0$,
- (b) $q = 4$ and $f_n \leq 0$ for all $n \geq 1$,

is satisfied, then the following implication is true. If

$$\lim_{n \rightarrow \infty} \mathbb{E}[I_q(f_n)^4] - 12\mathbb{E}[I_q(f_n)^3] = 12\nu^2 - 48\nu$$

then

$$\lim_{n \rightarrow \infty} \|f_n \star_r^\ell f_n\| = 0 \tag{5.14}$$

for all $r \in \{1, \dots, q\}$ and $\ell \in \{1, \dots, r \wedge (q-1)\}$ such that $(r, \ell) \neq (q/2, q/2)$,

$$\lim_{n \rightarrow \infty} \|f_n^2\| = 0, \tag{5.15}$$

$$\lim_{n \rightarrow \infty} \|f_n \tilde{\star}_{q/2}^{q/2} f_n - c_q f_n\| = 0 \quad \text{with } c_q = \frac{4}{(q/2)! \binom{q}{q/2}^2}. \tag{5.16}$$

Proof. First apply Lemma 5.6 to deduce that $A(I_q(f_n)) + R(I_q(f_n)) \rightarrow 0$, as $n \rightarrow \infty$. Assume that $q = 2$ and $\|f_n^2\| \rightarrow 0$. Then (5.15) is satisfied by assumption. Moreover,

$$R(I_2(f_n)) = 32\langle f_n \tilde{\star}_1^1 f_n, f_n \tilde{\star}_2^0 f_n \rangle - 48\langle f_n \tilde{\star}_2^0 f_n, f_n \rangle.$$

By the Cauchy-Schwarz inequality, we see that

$$|\langle f_n \tilde{\star}_1^1 f_n, f_n \tilde{\star}_2^0 f_n \rangle| \leq \|f_n \tilde{\star}_1^1 f_n\| \|f_n \tilde{\star}_2^0 f_n\|, \quad |\langle f_n \tilde{\star}_2^0 f_n, f_n \rangle| \leq \|f_n\| \|f_n \tilde{\star}_2^0 f_n\|.$$

With respect to the definition of the contractions, we see that $f_n \tilde{\star}_2^0 f_n = f_n^2$. We shall argue now that the sequence $\|f_n \tilde{\star}_1^1 f_n\|$ is bounded. For this, observe that for any fixed $(s, t) \in \mathcal{Z}^2$, we obtain by the Cauchy-Schwarz inequality that

$$\begin{aligned} |f_n \star_1^1 f_n(t, s)| &= \left| \int_{\mathcal{Z}} f_n(z, t) f_n(z, s) \mu(dz) \right| \\ &\leq \left(\int_{\mathcal{Z}} f_n^2(z, t) \mu(dz) \right)^{1/2} \left(\int_{\mathcal{Z}} f_n^2(z, s) \mu(dz) \right)^{1/2}. \end{aligned}$$

Consequently,

$$\|f_n \tilde{\star}_1^1 f_n\|^2 \leq \|f_n \star_1^1 f_n\|^2 = \int_{\mathcal{Z}^2} |f_n \star_1^1 f_n(t, s)|^2 \mu^2(d(s, t)) \leq \|f_n\|^4.$$

By assumption, we have that $\|f_n\|^2 \rightarrow \nu$, so the sequence is bounded. Now, the fact that $\|f_n^2\| \rightarrow 0$ implies that $R(I_2(f_n)) \rightarrow 0$. Hence, $A(I_2(f_n)) \rightarrow 0$, which implies that $A'(I_2(f_n)) \rightarrow 0$ using Lemma 5.7(1). Now, we apply Lemma 5.7(2) to see that (5.14) and (5.16) follow.

Next, let $q = 4$ and suppose that $f_n \leq 0$. Recall that the tensor product is bi-linear and it is easily verified that the contraction operation preserves this bi-linearity. Now, the fact that the kernels are non-positive ensures that $R(I_q(f_n)) \geq 0$ and we can again apply Lemma 5.7(1) to see that $0 \leq A'(I_q(f_n)) \leq A(I_q(f_n))$. Hence, we deduce that $A(I_q(f_n)) \rightarrow 0$. This directly implies that $A'(I_q(f_n)) \rightarrow 0$, such that the claim follows again by Lemma 5.7(2). \square

Remark 5.10. Let us explain in some more detail why in contrast to the case of normal approximation the kernels have to be non-positive for Gamma approximations. An inspection of the proof of Theorem 3.5 shows that a constant sign of the kernels is necessary to control the sign of scalar products. This is necessary in Lemma 5.4 and therefore also in Lemma 5.7 to control the signs of $A(I_q(f_n))$ and $A'(I_q(f_n))$, respectively. On the other hand, this is also necessary in part b) of Lemma 5.9, where one has to control the sign of $R(I_q(f_n))$. In this context, scalar products of the form $\langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle$, $p \in \{q/2 + 1, \dots, q\}$, appear. They are thrice-linear in f_n , such that $f_n \leq 0$ implies that $\langle f_n \tilde{\star}_p^{q-p} f_n, f_n \rangle \leq 0$ and we can conclude that $R(I_q(f_n)) \geq 0$. In summary, knowing that $A'(I_q(f_n)) \geq 0$ and $R(I_q(f_n)) \geq 0$ enables us to use part (2) of Lemma 5.7 to get the implication (ii) \implies (iii) in Theorem 3.5. Note that the latter scalar products in $R(I_q(f_n))$ actually stem from the third moment in assertion (ii) of Theorem 3.5 (see also (2.8)).

It is worth mentioning that this asymmetry in the assertions for Theorem 3.5 (and also in Theorem 3.3) is actually an intrinsic property of the Gamma distribution which contrasts the normal case. For the central limit theorem in a Poisson chaos, it can be easily seen that if the law of the sequence $\{I_q(f_n): n \geq 1\}$ converges to a standard normal law $\mathcal{N}(0, 1)$, then also the law $\{I_q(-f_n): n \geq 1\} = \{-I_q(f_n): n \geq 1\}$ converges to $\mathcal{N}(0, 1)$, since the standard normal law is symmetric. Consistently, assertions (ii) and (iii) in the four moments theorem for normal approximation are invariant under sign changes of the kernels. In sharp contrast, the assertions for Gamma approximations are not invariant under such sign changes, because of the lack of symmetry of the target distribution. This means that if the law of $\{I_q(f_n): n \geq 1\}$ converges to $\bar{\Gamma}_\nu$, then that law of $\{I_q(-f_n): n \geq 1\} = \{-I_q(f_n): n \geq 1\}$ cannot converge to $\bar{\Gamma}_\nu$. Consistently, assertions (ii) and (iii) in Theorem 3.5 inherit this asymmetry, which is reflected by the appearance of the third moment in (ii) and the term $\|f_n \tilde{\star}_{q/2}^{q/2} f_n - c_q f_n\|$ in (iii), both of them not being invariant under a change of the sign of f_n .

5.4. An alternative approach to the four moments theorem. In Remark 5.10 we explained that the sign condition on the kernels in part (b) of Theorem 3.5 ensures that $R(I_q(f_n)) \geq 0$. Together with $A'(I_q(f_n)) \geq 0$, this is sufficient in combination with part (2) of Lemma 5.7 to get the implication (ii) \implies (iii) in Theorem 3.5. On the other hand, for part (a) of Theorem 3.5, dealing with the case $q = 2$, the assumption that $\|f_n^2\| \rightarrow 0$ yields that $R(I_2(f_n)) \rightarrow 0$, an assertion also being sufficient in combination with $A'(I_q(f_n)) \geq 0$ to deduce the implication (ii) \implies (iii) in Theorem 3.5 from part (2) of Lemma 5.7. From this point of view, it is natural to ask whether the latter condition can be generalized to arbitrary

$q \geq 2$. Our next result shows that this is indeed possible, but leads to a result which is weaker than Theorem 3.5. Moreover, the proof again only works for $q = 4$ and we still have to impose a sign condition on the sequence of kernels.

Proposition 5.11. *Fix $\nu > 0$. Let $\{f_n: n \geq 1\} \subset L_s^2(\mu_n^4)$ be a sequence of kernels such that $f_n \geq 0$ for all $n \geq 1$ and such that the technical assumptions (A) and the normalization condition*

$$\lim_{n \rightarrow \infty} 4! \|f_n\|^2 = \lim_{n \rightarrow \infty} \mathbb{E}[I_4(f_n)^2] = 2\nu$$

are satisfied. Assume additionally that

$$\lim_{n \rightarrow \infty} \|f_n^2\| = 0, \text{ and } \lim_{n \rightarrow \infty} \|f_n \star_3^1 f_n\| = 0. \quad (5.17)$$

If the sequence $\{I_4(f_n)^4: n \geq 1\}$ is uniformly integrable, then the equivalence stated in Theorem 3.5 remains valid.

Proof. The implication (i) \implies (ii) follows from the uniform integrability of the sequence $\{I_4(f_n)^4: n \geq 1\}$ and (iii) \implies (i) is a consequence of Proposition 5.1. To establish the implication (ii) \implies (iii), we apply Lemma 5.6 and show that the term $R(I_4(f_n))$ defined at (5.9) converges to zero, as $n \rightarrow \infty$. With the Cauchy-Schwarz inequality we obtain for $p \in \{3, 4\}$ that

$$|\langle f_n \tilde{\star}_p^{4-p} f_n, f_n \rangle| \leq \|f_n \star_p^{4-p} f_n\| \|f_n\| \rightarrow 0,$$

since $\|f_n \star_p^{4-p} f_n\| \rightarrow 0$ and $\|f_n\|^2 \rightarrow \frac{\nu}{12}$. Moreover, for $p, r \in \{2, 3, 4\}$ with $p \neq r$ we also get

$$|\langle f_n \tilde{\star}_p^{4-p} f_n, f_n \tilde{\star}_r^{4-r} f_n \rangle| \leq \|f_n \tilde{\star}_p^{4-p} f_n\| \|f_n \tilde{\star}_r^{4-r} f_n\| \rightarrow 0.$$

The convergence is ensured by condition (5.17) if $p, r > 2$. If otherwise $p \wedge r = 2$, we use condition (5.17) together with the observation that $\|f_n \star_2^0 f_n\| = \|f_n \star_4^2 f_n\|$ and $\|f_n \star_2^2 f_n\| \leq \|f_n \star_4^4 f_n\|$ as a consequence of Fubini's theorem and the Cauchy-Schwarz inequality. Summarizing, we see that $R(I_4(f_n)) \rightarrow 0$, which in turn implies that $A(I_q(f_n)) \rightarrow 0$ thanks to Lemma 5.6. We can then conclude as in the proof of part (b) of Lemma 5.9. \square

Acknowledgements. We would like to thank Johanna F. Ziegel for initiating this collaboration and Giovanni Peccati for helpful discussions and valuable comments.

REFERENCES

- [1] S. Bourguin, Poisson convergence on the free Poisson algebra. Accepted for publication in Bernoulli (2014+).
- [2] P. de Jong, A central limit theorem for generalized quadratic forms. Probab. Theory Related Fields **75**, 261–277 (1987).
- [3] P. de Jong, A central limit theorem for generalized multilinear forms. J. Multivariate Anal. **34**, 275–289 (1990).
- [4] E.B. Dynkin and A. Mandelbaum, Symmetric statistics, Poisson point processes, and multiple Wiener integrals. Ann. Statist. **11**, 739–745 (1983).
- [5] P. Eichelsbacher and C. Thäle, New Berry-Esseen bounds for non-linear functionals of Poisson random measures. Electron. J. Probab. **19**, article 102 (2014).
- [6] R. Lachièze-Rey and G. Peccati, Fine Gaussian fluctuations on the Poisson space, I: contractions, cumulants and geometric random graphs. Electron. J. Probab. **18**, article 32 (2013).
- [7] G. Last, G. Peccati and M. Schulte, Normal approximation on Poisson spaces: Mehler's formula, second order Poincaré inequalities and stabilization. arXiv: 1401.7568 (2014).
- [8] G. Last, M.D. Penrose, M. Schulte and C. Thäle, Moments and central limit theorems for some multivariate Poisson functionals. Adv. in Appl. Probab. **46**, 348–364 (2014).

- [9] I. Nourdin and G. Peccati, Noncentral convergence of multiple integrals. *Ann. Probab.* **37**, 1412–1426 (2009).
- [10] I. Nourdin and G. Peccati, *Normal Approximations with Malliavin Calculus - From Stein's method to Universality*. Cambridge University Press, Cambridge (2012).
- [11] I. Nourdin, G. Peccati and G. Reinert, Invariance principles for homogeneous sums: universality of Gaussian Wiener chaos. *Ann. Probab.* **38**, 1947–1985 (2010).
- [12] D. Nualart and G. Peccati, Central limit theorems for sequences of multiple stochastic integrals. *Ann. Probab.* **33**, 177–193 (2005).
- [13] D. Nualart and J. Vives, Anticipative calculus for the Poisson process based on the Fock space. *Lecture Notes in Mathematics* **1426**, 154–165 (1990).
- [14] G. Peccati, J.L. Solé, M.S. Taqqu and F. Utzet, Stein's method and normal approximation of Poisson functionals. *Ann. Probab.* **38**, 443–478 (2010).
- [15] G. Peccati and M.S. Taqqu, Central limit theorems for double Poisson integrals. *Bernoulli* **14**, 791–821 (2008).
- [16] G. Peccati and M.S. Taqqu, *Wiener Chaos: Moments, Cumulants and Diagrams*. Bocconi and Springer, Milan (2011).
- [17] G. Peccati and C. Thäle, Gamma limits and U-statistics on the Poisson space. *ALEA Lat. Am. J. Probab. Math. Stat.* **10**, 525–560 (2013).
- [18] G. Peccati and C. Zheng, Multi-dimensional Gaussian fluctuations on the Poisson space. *Electron. J. Probab.* **15**, article 48 (2010).
- [19] G. Peccati and C. Zheng, Universal Gaussian fluctuations on the discrete Poisson chaos. *Bernoulli* **20**, 697–715 (2014).
- [20] M. Reitzner and M. Schulte, Central limit theorems for U -statistics of Poisson point processes. *Ann. Probab.* **41**, 3879–3909 (2013).
- [21] M. Schulte, Normal approximation of Poisson functionals in Kolmogorov distance. Accepted for publication in *J. Theor. Probab.* (2014+).
- [22] M. Schulte and C. Thäle, The scaling limit of Poisson-driven order statistics with applications in geometric probability. *Stoch. Proc. Appl.* **122**, 4096–4120 (2012).
- [23] M. Schulte and C. Thäle, Distances between Poisson k -flats. *Methodol. Comput. Appl. Probab.* **16**, 311–329 (2014).

TOBIAS FISSLER: INSTITUTE OF MATHEMATICAL STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF BERN, SWITZERLAND

E-mail address: tobias.fissler@stat.unibe.ch

CHRISTOPH THÄLE: FACULTY OF MATHEMATICS, RUHR UNIVERSITY BOCHUM, GERMANY

E-mail address: christoph.thaele@rub.de