Weathering versus circulation-controlled changes in radiogenic isotope tracer composition of the Labrador Sea and North Atlantic Deep Water

Friedhelm von Blanckenburg and Thomas F. Nägler

Isotopengeologie, Mineralogisch-Petrographisches Institut, Universität Bern, Bern, Switzerland

Abstract. Geological reconstructions and general circulation models suggest that the onset of both Northern Hemisphere glaciation, 2.7 Myr ago, and convection of Labrador Sea Water (LSW) were caused by the closure of the Panama Gateway ~ 4.5 Myr ago. Time series data that have been obtained from studies of ferromanganese crusts from the northwestern Atlantic suggest that radiogenic isotopes of intermediate ocean residence time (Pb and Nd) can serve as suitable tracers to reconstruct these events. However, it has been unclear until now as to whether the changes that have been observed in isotope composition at this time are the result of increased thermohaline circulation or due to the effects of increased glacial weathering. In this paper we adopt a box model approach to demonstrate that the shifts in radiogenic isotope compositions are unlikely to be due to changes in convection in LSW but can be explained in terms of increases of erosion levels due to the glaciation of Greenland and Canada. Furthermore, we provide experimental evidence for the incongruent release of a labile fraction of strongly radiogenic Pb and nonradiogenic Nd from continental detritus eroding into the Labrador Sea. This can be attributed to the glacial weathering of old continents and accounts for the paradox that one of the areas of the world most deficient in radiogenic Pb should provide such a rich supply of radiogenic Pb to the oceans. An important general conclusion is that the compositions of radiogenic isotopes in seawater are not always a reflection of their continental sources. Perhaps more importantly, the transition from chemical weathering to mechanical erosion is likely to result in significant variations in radiogenic tracers in seawater.

1. Introduction

Isotopic ocean tracers of intermediate residence time (Nd, Pb, Hf and Be) have been the subject of considerable attention in the past years in studies of long-term (Myr) [Burton et al., 1997; Christensen et al., 1997; Abouchami et al., 1999; Frank et al., 1999; Lee et al., 1999; von Blanckenburg and O'Nions, 1999; Piotrowski et al., 2000] and short-term (the past climate cycles [e.g., Rutberg et al., 2000] changes in ocean circulation and ocean chemistry. The reason for the interest in these tracers lies in the fact that their isotope ratios ¹⁴³Nd/¹⁴⁴Nd, ^{208,207,206}Pb/²⁰⁴Pb, ¹⁷⁶Hf/¹⁷⁷Hf, and ¹⁰Be/⁹Be are all distinct within different water masses. This is because their short to intermediate residence time limits the scale of homogenization and ensures a distinct signal from the sources of water masses. In particular, North Atlantic Deep Water (NADW) flowing south and Antarctic Bottom Water flowing north in the Atlantic Ocean are, in turn, labeled by distinctive Nd, Pb, Hf, and Be isotope ratios [Piepgras and Wasserburg, 1987; Kusakabe et al., 1990; von Blanckenburg et al., 1996; Albarède et al., 1998]. Furthermore, Nd isotopes allow for the detailed study of the evolution of Labrador Sea Water (LSW) through time as this water mass is characterized by a unique Nd isotope composition due largely to water from Baffin Bay [Stordal and Wasserburg, 1986], which is bordered by ancient continents (Figure 1).

Copyright 2001 by the American Geophysical Union.

Paper number 2000PA000550 0883-8305/01/2000PA000550\$12.00

This observation and the hypothesis that the closure of the Panama Gateway at 4.5 Ma might have been instrumental in initiating the Pleistocene mode of thermohaline circulation by increasing the amount of LSW flowing into NADW by convection [Stanley, 1995; Mikolajewicz and Crowley, 1997; Haug and Tiedemann, 1998] have led to extensive research efforts on isotope records on the Ma time frame. Such records of past isotope compositions in seawater can be reliably and quite easily obtained from the analysis of hydrogenetic ferromanganese crusts. North Atlantic Fe-Mn crusts show a marked decrease in ε_{Nd} (where ε_{Nd} is the ¹⁴³Nd/¹⁴⁴Nd ratio normalized to a chondritic value) from -11 to -13, an increase in ²⁰⁶Pb/²⁰⁴Pb from 18.8 to 19.2, and a decrease in both ²⁰⁷Pb/²⁰⁶Pb and ²⁰⁸Pb/²⁰⁶Pb over the past 3 Myr [Burton et al., 1997, 1999; Abouchami et al., 1999; Frank et al., 1999; Reynolds et al., 1999] (see also Figure 2). These Nd and Pb isotope changes would be compatible with a gradual strengthening of Labrador Sea Water convection. However, similar isotope changes could be accounted for by increased erosion, namely, an increase in the dominance of mechanical erosion versus chemical weathering in Greenland and North America, arising from the Northern Hemisphere glaciation and leading to similar shifts [Burton et al., 1997, 1999; von Blanckenburg and O'Nions, 1999]. One of the difficulties inherent in the use of radiogenic tracers in paleoceanography is the fact that both changes in ocean circulation and changes in weathering of the tracer's sources might lead to secular variations in the isotope ratios. Thus it will often not be possible to single out the driving forces that determine isotopic changes [von Blanckenburg, 1999].



Figure 1. Circulation of deep water in the North Atlantic (solid arrow) and surface circulation through Baffin Bay (dashed arrow). Numbers in ellipses give the amount of deep water formed at these sites in sverdrup ($1 \text{ Sv} = 1 \times 10^6 \text{ m}^3 \text{ s}^{-1}$). A total of 18 Sv of North Atlantic Deep Water (NADW) are formed by convection in the Greenland-Iceland-Norwegian Seas (11 Sv) and in the Labrador Sea (7 Sv [Schmitz and McCartney, 1993; Dickson and Brown, 1994]. Boxed numbers in italic type are $^{206}\text{Pb}/^{204}\text{Pb}$ ratios; other numbers are ϵ_{Nd} of both source rocks [Moorbath et al., 1997; Winter et al., 1997] and Pb and Nd dissolved in seawater [Stordal and Wasserburg, 1986; Piepgras and Wasserburg, 1987; von Blanckenburg et al., 1996]. Note that Baffin Bay is surrounded by ancient continents where ϵ_{Nd} is as low as -60 [Moorbath et al., 1997]. In areas surrounding the Greenland-Iceland-Norwegian Seas, ϵ_{Nd} is -20 to -8 (compilation by Winter et al. [1997]. Also shown are the sampling sites of ODP Hole 645 (Baffin Bay) and NW Atlantic Fe-Mn crusts ALV 539, BM 1965.05 [Burton et al., 1997; O'Nions et al., 1998; von Blanckenburg and O'Nions, 1999], Blake, and Antilles [Reynolds et al., 1999].

The aim of this paper is to shed light on the relative importance of weathering effects versus changes in thermohaline circulation for variations in the isotope composition of NADW in the Pleistocene. As a first step we tested the sensitivity with which the isotope composition responds to convection of LSW by utilizing box models of tracer exchange between Baffin Bay and NADW. We will demonstrate that the composition of radiogenic tracers in the North Atlantic responds sensitively to changes in the style of weathering and is insensitive to changes in the strength of thermohaline circulation. To further test the possibility that the causes are changes in the style and degree of weathering, we have performed leaching experiments on the materials that constitute the sources of tracers that erode into the Labrador Sea. These experiments simulate the chemical release of tracers due to mechanical erosion of materials, such as those that prevail in strongly glaciated areas.

2. Experimental Approaches and Results

2.1 Tracer Exchange by Changes in the Thermohaline Circulation

The gradual increase of $^{206}Pb/^{204}Pb$ from 18.9 to 19.2 and the concomitant decrease of ε_{Nd} in the NW Atlantic from -11 to -13 in the past 3 Myr (Figure 2) might be generated by strengthening of LSW convection. This is a possibility discussed frequently in the literature [*Burton et al.*, 1997; *Heinze*



Figure 2. (a) The ²⁰⁶Pb/²⁰⁴Pb isotope ratio variations and (b) ε_{Nd} variations in northwest Atlantic Deep Water as recorded by hydrogenetic Fe-Mn crusts [*Burton et al.*, 1999; *O'Nions et al.*, 1998; *Reynolds et al.*, 1999]. These crusts have been dated using the radioactive decay of cosmogenic ¹⁰Be, which reduces their ¹⁰Be/⁹Be ratio with time. Note that the major increase in ²⁰⁶Pb/²⁰⁴Pb and decrease in ε_{Nd} are after 3 Myr, the onset of Northern Hemisphere glaciation, whereas the major hydrographic changes attributed to the closure of the Panama Gateway have taken place at 4.5 Ma [*Haug and Tiedemann*, 1998].

and Crowley, 1997; Abouchami et al., 1999; Burton et al., 1999; Vance and Burton, 1999]. In order to test this hypothesis the exchange of tracers between Baffin Bay and the entire North Atlantic due to advection was simulated in a simple two-box model (Figure 3). First, an increase of LSW convection was simulated by varying the exchange between Baffin Bay and the North Atlantic between 0 and 10 Sverdrup (1 Sv = 1×10^6 m³ s⁻¹). Resulting ε_{Nd} in the North Atlantic are shown in Figure 3a. It is noteworthy that resulting ratios are constant at -10.5 for equal Nd fluxes per unit area into both basins and at -13 for a tenfold increase in Nd flux into Baffin Bay. With the exception of very low exchange rates and a very high Nd flux into Baffin Bay, isotope ratios are virtually independent of the variations in water exchange. This result becomes more obvious when considering the fact that at a present-day exchange rate of 2 Sv and with the volume given in Figure 3, the contents of Baffin Bay are replenished every 15 years. The Labrador Sea, with a much larger volume, is also replenished over similar timescales because the deep water formation rate is also higher at 7 Sv. In this case the water replenishment rate is much shorter than the tracer's residence time τ , and this tracer's budget in the Atlantic turns out to be insensitive to the rate of water exchange with LSW. Therefore intermediate- τ tracer advection from one box to the other is essentially limited by the rate at which tracers enter the basin through erosion, regardless of the actual water exchange rate. The only exception to this prediction is if exchange rates are only slightly higher than a total shutoff of LSW convection. In this case the water replenishment time approaches the tracer's residence time, and the North Atlantic's Nd composition becomes sensitive to the water exchange rate with LSW (Figure 3a). The Nd isotope composition of Baffin Bay water resulting from the same experiment (not shown) reacts sensitively to any change in water exchange flux, and the sense of the change is opposite to that in the North Atlantic. Baffin Bay is where changes in deep water generation would show up in the Nd isotope record, whereas in the North Atlantic they would not. However, suitable Fe-Mn crusts have not been recovered from Baffin Bay.

The same simulation has been performed for Pb. Since Pb has a North Atlantic Deep Water residence time τ of 40 years [Cochran et al., 1990], which is much closer to the water replenishment time of Baffin Bay or Labrador Sea, Pb isotope variations would be detectable both in the North Atlantic and in Baffin Bay, in particular if the flux of Pb per unit area into Baffin Bay is much higher than that into the Atlantic (Figure 3b). However, the sense of a shift resulting from strengthening of LSW convection, which is governed by the low Pb isotope composition of the areas surrounding the Labrador Sea (Figure1), is opposite to that observed in the NW Atlantic Fe-Mn crust records (Figure 2a). Therefore, at first sight, the observed Pb isotope compositions are also incompatible with a gradual strengthening of LSW convection. The only scenario that is compatible with the observed Pb isotope trends would be a combination of the strenghtening LSW convection, a high terrigenous Pb flux into Baffin Bay, and an incongruent release of radiogenic Pb, as explored in section 2.2.

Baffin Bay Water has a distinctly low Nd isotope composition (Figure 1). Baffin Bay is also the basin with the highest dissolved Nd concentrations of any basins worldwide [*Stordal* and Wasserburg, 1986]. These observations make a high terrigenous flux of Nd into the basin likely. In Figure 3c the flux of Nd into Baffin Bay was increased up to tenfold relative to that of the North Atlantic per unit area while leaving the water exchange rate constant. A strong decrease in North Atlantic ε_{Nd} results. An approximately fivefold increase in erosional input would be sufficient to explain the Pleistocene shift in isotope ratios measured in NW Atlantic Fe-Mn crusts. It must be noted, however, that such an increase must be particularly strong in the Labrador Sea area, as a simultaneous increase of Nd released into the other sources of NADW would result in constant isotope compositions. This is not entirely unlikely,



Figure 3. Isotopic ratios calculated from two-box models of tracer exchange between Baffin Bay and the North Atlantic. Results shown are those for the North Atlantic box. The shaded areas are the compositions of NW Atlantic Fe-Mn crusts at different times before and after closure of the Panama Gateway. The Atlantic box has a volume of 1.69 × 10¹⁷ m³; Baffin Bay has a volume of 9.6×10^{14} m³. Tracer transport is by advective exchange between Baffin Bay and the North Atlantic. Today this is ~2 Sv [Schmitz and McCartney, 1993; Dickson and Brown, 1994]. This rate is used for the standard model unless indicated otherwise. Tracers are present for which the removal by scavenging is governed by their residence time, which is 40 years for Pb, 250 years for Be, and 1000 years for Nd. Erosional tracer flux (total Pb, Nd and ⁹Be) is proportional to the area of the boxes and is 1 mass unit cm⁻² yr⁻¹ in the standard model and 10 mass units cm⁻² yr⁻¹ into Baffin Bay for the models of Figures 3a and 3b, shown by squares. Atmospheric ¹⁰Be deposition is also uniform over the area and has been held constant at 0.4 mass units cm⁻² yr⁻¹, such that an "Atlantic-type" $^{10}Be^{9}Be$ ratio of 0.4 x 10⁻⁷ results. The input of Nd into the North Atlantic box is characterized by an ε_{Nd} of -10, and that into Baffin Bay is characterized by -40. In Figure 3c a model is added where the ENd of the input to Baffin Bay is -30 [Moorbath et al., 1997; Piepgras and Wasserburg, 1987; Stordal and Wasserburg, 1986]. Input 206 Pb/204 Pb is 19 in the North Atlantic and 16 in Baffin Bay. It must be noted that the conclusions are not affected by the choices made for the input isotope ratios; it is the relative changes in ratios that matter. (a and b) The effect of strengthening of LSW, which was simulated by varying the water exchange rate between Baffin Bay and the North Atlantic between 0 and 10 Sv for two different Nd and Pb fluxes into Baffin Bay. (c) The effect of an increased glacial erosional input into Baffin Bay on North Atlantic Nd chemistry, which was simulated by increasing the flux of Nd up to tenfold for two different ε_{Nd} of the input into Baffin Bay. (d) The effect of an increased glacial erosional input into Baffin Bay for the North Atlantic ¹⁰Be/⁹Be ratio, which was simulated by increasing the flux of ⁹Be into Baffin Bay up to tenfold while leaving the ¹⁰Be flux constant. Runs were also conducted for an exchange between the entire Labrador Sea and the North Atlantic. Because the higher volume of the Labrador Sea is matched by higher water exchange rates, results are very similar to those of the Baffin Bay model.

given the low ratio of the Laboratory Sea's area to the length of its margins, when compared with the other North Atlantic basins.

It has been argued that an increase in erosional flux into the Labrador Sea is an unlikely cause for radiogenic tracer variations, as it would be accompanied by ⁹Be, which would lower the ${}^{10}\text{Be}/{}^9\text{Be}$ ratio in the same period [Burton et al., 1999]. This is apparently in contradiction to the observation that initial ${}^{10}\text{Be}/{}^9\text{Be}$ ratios in the NW Atlantic have been constant over the past 7 Myr [von Blanckenburg and O'Nions, 1999]. It can be demonstrated with a fourth model, however, that relative to the entire North Atlantic, the overall increase in the

Table 1. Pb Isotope Results of Le	each Experiments ^a									
Sample	²⁰⁸ Pb/ ²⁰⁴ Pb	²⁰⁷ Pb/ ²⁰⁴ Pb		²⁰⁶ Pb/ ²⁰⁴ Pb		²⁰⁸ Pb/ ²⁰⁶ Pb		²⁰⁷ Pb/ ²⁰⁶ P	۹ I	Pb, ppm
Core B Hole 1X, 2.4 mbsf Leach 0.05 M HCl	38.262 ± 0.020 43.254 ± 0.113	Baffin Ba 15.3789 ± (15.5686 ± (y <i>ODP Leg</i> 0.0080 0.0401	$\begin{array}{rcrcr} 105 \ Core \ 645 \ Sed \\ 17.3151 \ \pm \ 0.0 \\ 20.4461 \ \pm \ 0.0 \end{array}$	iments 086 534	2.20973 ± (2.11549 ± ().00024).00049	0.88818 ± 0.76144 ±	0.00010 0.00019	5.7 0.04
Core B Hole 5X, 35 mbsf Leach 0.05 M HCl	38.152 ± 0.018 42.044 ± 0.116	15.3837 ± 15.7483 ±	0.0071 0.0409	17.1386 ± 0.0 21.0222 ± 0.0	079 558	2.22608 ± (1.99997 ± ().00025).00052	0.89760 ± 0.74913 ±	0.00010 0.00019	7.6 0.06
Core B Hole 8X, 64 mbsf Leach 0.05 M HCl	36.901 ± 0.016 42.701 ± 0.154	15.2377 ± 15.7122 ±	0.0066 0.0566	16.3114 ± 0.0 20.5099 ± 0.0	070 760	2.26227 ± (2.08196 ± (0.00025 0.00069	0.93417 ± 0.76608 ±	0.00010 0.00030	8.0 0.02
Core B Hole 9X, 76 mbsf Leach 0.05 M HCl	39.445 ± 0.016 42.008 ± 0.031	15.6162 ± 15.8508 ±	0.0063 0.0104	19.0946 ± 0.0 21.2244 ± 0.0	077 132	2.06579 ± (1.97922 ± (0.00022 0.00037	0.81783 ± 0.74682 ±	0.00008	90 1.1
Core B Hole 11X, 94 mbsf Leach 0.05 M HCl	36.461 ± 0.021 44.916 ± 0.048	15.2944 ± 15.8366 ± 0	0.0090	16.7009 ± 0.0 21.9121 ± 0.0	099 229	2.18320 ± (2.04984 ± (0.00024 0.00031	0.91578 ± 0.72273 ±	0.00001	2.9 0.1
SM-AC-95-9 Acasta Gneiss Leach 0.05 M HCi	35.323 ± 0.017 42.775 ± 0.198	15.2228 ± 15.3439 ±	Archu 0.0073 0.0711	ean Rocks 15.2510 ± 0.0 18.9061 ± 0.0	071 899	2.31614 ± (2.26250 ± ().00024).00074	0.99815 ± 0.81158 ±	0.00010 0.00043	4.1 0.02
SM-AC-95-11 Acasta Gneiss Leach 0.05 M HCI	$\begin{array}{rrrr} 43.178 \pm 0.019 \\ 127.792 \pm 0.096 \end{array}$	15.4833 ± 16.6784 ±	0.0068 0.0125	$15.5226 \pm 0.0 \\ 22.1970 \pm 0.0 \\ \end{array}$	068 166	2.78160 ± (5.75717 ± ().00029).00083	0.99747 ± 0.75138 ±	0.00010 0.00009	6.3 0.5
SM-AC-95-12 Acasta Gneiss Leach 0.05 M HCI	$\begin{array}{rrrr} 43.125 \pm 0.018 \\ 48.117 \pm 0.023 \end{array}$	16.3038 ± 16.1880 ±	0.0068 0.0071	18.6414 ± 0.0 18.9831 ± 0.0	078 080	2.31339 ± (2.53470 ± (0.00025 0.00042	0.87460 ± 0.85276 ±	0100000	14.6 3.2
155710 Diorite, Amîtsoq Gneiss Leach 0.05 M HCl	31.510 ± 0.013 31.979 ± 0.023	13.1985 ± 13.3310 ±	0.0054 0.0096	11.5276 ± 0.0 12.1225 ± 0.0	047 083	2.73341 ± (2.63797 ± ().00028).00043	1.14495 ± 1.09969 ±	0.00012 0.00016	7.2 0.1
110528 65°06,76N, 49°56,37W Leach 0.05 M HCl	33.448 ± 0.014 39.776 ± 0.019	Amii 14.1312 ± 0 15.2644 ± 0	tsoq and Isu 0.0057 0.0069	a Stream Sedimen 13.6887 ± 0.0 19.4698 ± 0.0	<i>ts</i> 055 086	2.44349 ± (2.04296 ± ().00026).00032	1.03232 ± 0.78400 ±	0.00010 0.00009	66.0 0.6
110326 65°09,77N, 49°47,83W Leach 0.05 M HCl	33.422 ± 0.014 39.279 ± 0.019	14.2131 ± 0 14.9146 ± 0	0.0057 0.0068	14.0472 ± 0.0 19.2477 ± 0.0	056 085	2.37925 ± (2.04069 ± ().00025).00029	1.01181 ± 0.77488 ±	0.00010 0.00008	70 0.5
105862 65°09,84N, 50°07,01W Leach 0.05 M HCl	34.361 ± 0.014 36.859 ± 0.016	14.4182 ± (14.8489 ± (0.0058 0.0065	14.8197 ± 0.00 16.7815 ± 0.00	090 069	2.31861 ± (2.19642 ± ().00024).00027	0.97291 ± 0.88484 ±	0.00010 0.00010	112 1.3
105894 65°08,04N, 49°52,79W Leach 0.05 M HCI	33.222 ± 0.014 39.642 ± 0.020	14.0779 ± (14.7519 ± (0.0057 0.0068	13.4561 ± 0.00 17.8670 ± 0.00	054 078	2.46891 ± (2.21874 ± ().00027).00039	1.04621 ± 0.82565 ±	0.00011 0.00010	119 0.6

VON BLANCKENBURG AND NÄGLER: RADIOGENIC OCEAN TRACERS

_
\sim
ີ
_
_
_
.=
-723
_
_
0
-
u
2
9
9
9
С
1 . (c)
1. (c
e 1. (c
le 1. (c
ole 1. (c
ble 1. (c
able 1. (c
able 1. (c

Sample	²⁰⁸ Pb/ ²⁰⁴ Pb	²⁰⁷ Pb/ ²⁰⁴ F	ą	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁸ Pb/ ²⁰⁶ Pb	²⁰⁷ Pb/ ²⁰⁵ Pb	Pb, ppm
		Amitsoq	and Isua Str	ream Sediments (contin	ed)		
110356 65°04.97N. 50°08.29W	33.334 ± 0.014	14.0953 ±	0.0057	13.9307 ± 0.0056	2.39286 ± 0.00026	1.01182 ± 0.0001	0 80
Leach 0.05 M HCl	36.672 ± 0.030	14.5998 ±	0.0101	16.4439 ± 0.0109	2.23013 ± 0.00113	0.88786 ± 0.0003	0 0.4
			Multip	le Leach Test			
Core E Hole 4R 456 mbsf	35.832 ± 0.017	15.1885 ±	0.0074	15.8756 ± 0.0077	2.25703 ± 0.00025	0.95672 ± 0.0001	0 8.2
0.05 M HCI	40.706 ± 0.058	15.7733 ±	0.0221	21.0507 ± 0.0304	1.93370 ± 0.00037	0.74930 ± 0.000	2 0.1
0.6 M HCI	40.887 ± 0.029	15.7287 ±	0.0103	20.4993 ± 0.0129	1.99454 ± 0.00044	0.76728 ± 0.0001	4 0.5
0.05 M HBr	40.767 ± 0.073	15.6943 ±	0.0267	20.7567 ± 0.0336	1.96405 ± 0.00067	0.75611 ± 0.0002	0 0.2
NaCl + 0.05 M HBr pH=1.5	40.626 ± 0.080	15.6851 ±	0.0297	20.2258 ± 0.0389	2.00863 ± 0.00062	0.77550 ± 0.0002	2 0.1
^a An amount of 0.5 g of the detrita	I fraction of Pleistocene (DDP 645 sedim	ents [Srivas	tava et al., 1989], crusl	ed Archean rocks [Moorbath et a	L, 1997], and Greenland	iver sand was

MS) using a Tl doping technique [*Belshaw et al.*, 1999]. Known amounts of Tl were also used to measure approximate Pb concentrations. Comparison with isotope dilution shows that this is mostly accurate to within a factor of two. Repeated analysis of NBS 981 yielded ²⁰⁸Pb²⁰⁴Pb = 36.6944 ± 440 ppm, ²⁰⁷Pb²⁰⁴Pb = 15.4888 ± 400 ppm, ²⁰⁶Pb²⁰⁴Pb = 16.9334 ± 420 ppm, ^{20§}Pb/²⁰⁶Pb = 2.1670 \pm 60 ppm, and ²⁰⁷Pb/²⁰⁶Pb = 0.9147 \pm 120 ppm (2 σ standard deviation, n = 9). Isotope ratios in Table 1 are presented as measured, but quoted errors include to avoid decomposition of apatite or other accessory minerals. One sample was also treated with stronger HCl, mild HBr, and a NaCl-HBr solution with seawater salinity. Results were similar for all leach proce-From mass balance considerations it follows that the leach residues are not resolvably different in isotope composition exchange. Pb was measured by a Nu-Instruments multicollector inductively coupled plasma-mass spectrometer (ICPand Harlavan et al. [1998] seafloor below er used by Erel meters sample runs with the external standard deviation of standards. Here mbsf is This leach was much milder than mild agitation. by HF. purified by HNO₃-HBr anion For comparison, 50 mg of bulk rock was decomposed 24 hours under exposed to 5 mL of cold 0.05 M HCl for precision of Samples were propagation of the internal total materials. to the I dures.

⁹Be flux associated with an increase in erosion around Baffin Bay is so small that a virtually unresolvable shift in North Atlantic ¹⁰Be/⁹Be would result (Figure 3d). This is because even a large increase in the flux of ⁹Be into the Labrador Sea results in only a minute shift in the total ⁹Be budget in the Atlantic.

2.2. Tracer Release Experiments

As noted above, the observed increase in the North Atlantic's radiogenic Pb over the last 3 Myr is incompatible with either an increase in LSW convection or an overall increase in erosion of the Labrador and Greenland areas. This is because these areas have a Pb isotope composition opposite of what would be required to drive this shift (Figure 1). Our explanation for this paradox is that rocks undergoing physical erosion, such as in glaciated areas, will incongruently release a radiogenic fraction of Pb or Sr [Erel et al., 1994; Harlavan et al., 1998]. In contrast, soils that have experienced prolonged weathering release tracers having the isotope composition of the bulk soil. Similar observations have been made for Nd: around 70% of the Nd is lost from the topsoil of weathered tills, and the residual ¹⁴³Nd/¹⁴⁴Nd in the soil was higher by up to 4 ε_{Nd} units than in the unweathered C horizon [Öhlander et al., 2000]. In a boreal river, ε_{Nd} was lower by almost 10 ε_{Nd} units than in the unweathered bedrock in the area [Andersson et al., 2001]. This indicates the release of a strongly nonradiogenic Nd fraction into solution from physically eroded materials.

To investigate the tracer release from the weathering of rocks in the areas surrounding the Labrador Sea, Archean rocks, river sediments from Archean Greenland, and the detrital fraction of Pleistocene Ocean Drilling Program sediments from Baffin Bay have been treated using mild acid leaches which have been measured for their Pb and Nd isotope composition and compared to the original bulk material (Tables 1 and 2). The analytical details are given in Table 1.

Results (Figure 4) show that in all cases a labile pool of Pb is available that is much more radiogenic than the bulk rock. For example, a total sediment has a $^{\overline{206}Pb/^{204}Pb}$ of 13, with a leacheate ²⁰⁶Pb/²⁰⁴Pb of 19. Similar effects are also visible for ²⁰⁸Pb/²⁰⁶Pb and ²⁰⁷Pb/²⁰⁶Pb. Multiple leach tests further show that this radiogenic fraction is always released, regardless of the reagent used (Table 1). These experiments obviously do not precisely simulate nature in terms of timescale and solution chemistry. However, the simple fact that this labile radiogenic Pb is available for release in various aqueous solutions makes it very likely that this fraction would be the first to be released into seawater, too. The amount of radiogenic Pb lost from the deposited sediments is small. Therefore the detrital sediments will, on the whole, retain their original isotope composition, while the overlying water column has a much more radiogenic composition. The sense of these Pb isotope shifts are exactly the same as those experienced by NADW in the period from 7 Ma to the present (Figure 5). Therefore the variations in NADW Pb isotope composition are compatible with an increasing amount of mechanically eroded material introduced into Baffin Bay and the Labrador Sea.

Sample	Nd, ppm	Sm ^b , ppm	¹⁴⁷ Sm/ ¹⁴⁴ Nd	¹⁴³ Nd/ ¹⁴⁴ Nd	ε _{Nd} (0)
Ba	iffin Bay ODP Le	g 105 Core 64	5 Sediments		
Core B Hole 1X, 2.4 mbsf	6.37	1.20	0.1135	0.511492 ± 27 ^c	-22.4
Leach 0.05 M HCl	0.189	0.0362	0.1161	0.511227 ± 18°	-27.5
Core B Hole 5X, 35 mbsf	7.72	1.32	0.1033	0.511242 ± 20^{a}	-27.2
Leach 0.05 M HCl	0.200	0.004	0.1247	0.511609 ± 72^{d}	-20.1
Core B Hole 8X, 64 mbsf	67.28	12.30	0.1105	0.511141 ± 20^{a}	-29.2
Leach 0.05 M HCl	0.109	0.0190	0.1054	0.511208 ± 19^{a}	-27.9
Core D Hole 14R, 400 mbsf	4.35	0.792	0.1101	$0.511446 \pm 27^{\circ}$	-23.3
Leach 0 05 <i>M</i> HC1	0.545	0.110	0.1217	0.511745 ± 27°	-17.4
Core E Hole 4R, 456 mbsf	7.27	1.32	0.1094	0.511452 ± 28^{a}	-23.1
Leach 0.05 M HC1	1.30	0.273	0.1272	0.511961 ± 22^{a}	-13.2
Core D Hole 9R, 504 mbsf	7.60	1.39	0.1107	0.511476 ± 96 [°]	-22.7
Leach 0.05 M HCl	1.18	0.247	0.1267	0.511607 ± 25 [°]	-20.1
	Amitsog and Is	ua Stream Sea	diments		
110528 65°06,76N, 49°56,37W	54.26	11.42	0.1272	0.510947 ± 22^{a}	-33.0
Leach 0.05 M HC]	1.11	0.180	0.0984	0.510358 ± 31°	-44.5
110326 65°09,77N, 49°47,83W	45.7	11.3	0.1490	0.511397 ± 26°	-24.2
Leach 0.05 M HCl	1.14	0.215	0.1133	$0.510541 \pm 20^{\circ}$	-40.9
105862 65°09,84N, 50°07,01W	82.7	15.8	0.1151	0.5110881 ± 22 ^a	-34.3
Leach 0.05 M HCl	3.36	0.534	0.0959	$0.510401 \pm 32^{\circ}$	-43.6
105894 65°08,04N, 49°52.79W	39.9	8.90	0.1347	0.511302 ± 32^{a}	-26.1
Leach 0.05 M HCl	1.53	0.223	0.0884	0.510500 ± 28^{d}	-41.7

Table 2. Nd Isotope Results of Leach Experiments^a

^a Mass balance calculations show that leach residues (not measured) are all within 0.7 ε_{Nd} units to those of the corresponding totals.

^b Sm concentrations were measured by isotope dilution on MC-ICP-MS.

^c Analyses were measured by Thermal Ionization Mass Spectrometry (TIMS) as NdO+ on a modified AVCO single collector instrument (¹⁴³Nd/¹⁴⁴Nd external reproducibility is 0.000034, 2σ standard deviation, n = 5). The TIMS results were adjusted by 1 ε_{Nd} for instrument bias to adjust for a La Jolla value of 0.511860.

^d Samples were measured on a Nu-Instruments multicollector ICP-MS (143 Nd/ 144 Nd external reproducibility is 0.000019, 2 σ standard deviation, n = 15). No corrections were required for the Nu ICP-MS data.

Fresh Greenland river sediments release a significantly nonradiogenic fraction of Nd (Table 2 and Figure 6). This fraction amounts to 2-3% of the bulk rock's Nd and is between 10 and 15 ε_{Nd} units less radiogenic than the bulk rock. Leacheates of Baffin Bay ODP sediments show smaller differences with respect to their bulk material than those of fresh river sediments, and mostly, the reverse systematics in both ε_{Nd} and the Sm/Nd ratio are seen (Table 2 and Figure 6). We consider the more radiogenic Nd fraction measured in the leacheate of ODP sediments to be dominated to a large extent by the fraction that has been adsorbed from seawater. Such Nd would be present in relatively high concentrations in the Fe-Mn oxyhydroxide surface coatings of the minerals [*Vance and Burton*, 1999] and would reflect the isotope composition of the overlying water column. In surface LSW 2.4 Myr ago ε_{Nd} was -13, and decreased to -18 in LSW and to -22 in Baffin Bay today [*Stordal and Wasserburg*, 1986; *Vance and Burton*, 1999]. The increase in ¹⁴⁷Sm/¹⁴⁴Nd in the older leacheates is also compatible with a shift from high Sm/Nd ratios in seawater 2.4 Myr ago to low ratios observed today [*Vance and Burton*, 1999]. These shifts in LSW Nd isotope composition and Sm/Nd ratio reflect the increased input of labile low ε_{Nd} and low Sm/Nd after the onset of Northern Hemisphere glaciation released from mechanically eroded glacial tills.

The question thus arises as to the availability of radiogenic Pb in ODP sediments, while at the same time, nonradiogenic Nd fraction in the same materials is absent. This contrasts with the behavior of fresh river sediments, which show a greatly incongruent release of both radiogenic Pb and nonradiogenic Nd. Such differential behavior might be due to



Figure 4. Pb isotope ratios of dissolved bulk samples (solid squares) and 0.05 M HCl leacheates (open squares). The latter represent the Pb available for release into seawater. Analytical details are given in Table 1. Analytical errors are smaller than symbol sizes.

the fact that the labile nonradiogenic Nd is immediately lost upon the immersion of the sediment in seawater, whereas a large fraction of the radiogenic Pb is still available for release. This suggestion is compatible with the large fraction of Nd (70%) lost during weathering of glacial tills [*Öhlander et al.*, 2000].

The incongruent release of radiogenic Pb and of nonradiogenic Nd is typical for a strong mechanical erosion regime where an abundance of fresh mineral surfaces have been opened for weathering. Evidence for weathering of mechanically eroded materials is provided in soils and clastic sediment by the availability of a labile radiogenic pool of ²⁰⁸Pb and ²⁰⁶Pb, which is predominantly hosted by accessory minerals (mainly monazite, allanite and xenotime) which might have experienced radiation damage [*Erel et al.*, 1994], or by grain boundaries rich in the parent isotopes ²³⁸U and ²³²Th [*Silver et al.*, 1984]. The low ε_{Nd} and the low Sm/Nd ratio point to accessory minerals enriched by light rare earth element or grain boundaries as hosts for this labile Nd fraction, too. Upon prolonged chemical weathering this pool is



Figure 5. Pb isotope ratio shifts of NADW as recorded in northwest Atlantic Fe-Mn crusts [O'Nions et al., 1998; Burton et al., 1999; Reynolds et al., 1999]. Note that these shifts have the same sense as those between bulk rock and leacheates of the experiments shown in Figure 2.

lost into solution [Erel et al., 1994; Harlavan et al., 1998; Öhlander et al., 2000; Andersson et al., 2001]. This mechanism resolves the contradiction that the Labrador Sea provides the most radiogenic seawater Pb from an area where source rocks are among the least radiogenic in the world (Figure 1). A strong mechanical erosion regime is in operation on the North American and Greenland Shields, where glacial tills are the predominant sources for marine detritus. In support of this the differences in all Pb and Nd isotope ratios between the labile fraction and the bulk stream sediment follow the same trend as the early Pleistocene to recent shifts seen in the northwest Atlantic Fe-Mn crusts (Figures 2 and 4). These shifts can thus be attributed to a gradual shift from prolonged chemical weathering to mainly physical erosion during the buildup of the Northern Hemisphere ice sheets. Such incongruent release of isotope tracers would reduce the flux of bulk terrigenous material that otherwise would be required to satisfy the Labrador Sea Water's extreme isotope compositions (section 2.1).

3. Discussion and Conclusions

The obvious conclusion is that NADW Nd isotope compositions are insensitive to changes in the LSW contribution. This is in accordance with the suggestion that Nd in LSW itself underwent a major change toward less radiogenic compositions after 2.5 Ma [Vance and Burton, 1999]. Therefore the changes reported in NW Atlantic seawater isotope composition after the onset of Northern Hemisphere glaciation are best interpreted as a consequence of shifts in the style of weathering from chemical to more mechanical (glacial) erosion and also as due to an overall increase in the amount of erosion. Both are related to the formation of ice sheets in Greenland and the Canadian Shield with an increasing supply of icerafted debris [Larsen et al., 1994]. Incongruent weathering of areas surrounding the other sources of NADW, such as the Fennoscandian crust or eastern Greenland, would result in isotope shifts of the same sense as those observed for the Labrador Sea. It has been observed that glacial weathering in the areas surrounding the Arctic seas results in the release of radiogenic Pb, nonradiogenic Nd, and radiogenic Sr [Winter et al., 1997].

Other radiogenic isotope seawater tracers might be subject to the same mechanism: Hafnium isotopes from the same northwest Atlantic Fe-Mn crusts as those presented in Figure 2 show a shift to less radiogenic values (ε_{Hf} shifts from +3 to – 1 in the past 4 Myr) that virtually mirrors that of Nd in the same crusts [*Piotrowski et al.*, 2000]. This is compatible with the exposure of freshly eroded zircon minerals by physical erosion. These are rich in nonradiogenic Hf. Furthermore, the gradual increase in global seawater ¹⁸⁷Os/¹⁸⁸Os in the Quaternary was explained by progressive removal of a deeply weathered pre-Quaternary regolith from the Precambrian shields surrounding the Labrador Sea [*Peucker-Ehrenbrink and Blum*, 1998].

It must be pointed out that some reorganization of smallscale ocean circulation might be visible in the Fe-Mn crust records. For example, the westward shift with time of a local water mass rich in radiogenic Pb might be responsible for the increase in radiogenic Pb of the Fe-Mn crust ALV 539 (Figures 1 and 2). This shift precedes that experienced by the more westerly crusts by ~1 Ma. In a similar way, a time lag between the decrease in ε_{Nd} between crust Alv 539 and BM 1965.05 has been explained by vertical shifts of water masses [Burton et al., 1999]. It must be noted, however, that the apparent difference between the Nd and Pb isotope time series from ALV 539 and other NW Atlantic crusts has been attributed to a possibly inconsistent chronology [Reynolds et al., 1999]. The overall North Atlantic shifts in Pb, Nd, and Hf isotope compositions in the past 3-4 Myr must, however, be caused by changes in erosion rather than lateral reorganization of local circulation: The shifts are visible over large areas, from the east Atlantic [Abouchami et al., 1999] to the Carribean [Reynolds et al., 1999]; they affect tracers of entirely



different deep-ocean τ (Pb, Nd, Hf); and they are absent in initial ¹⁰Be/⁹Be [von Blanckenburg and O'Nions, 1999].

The results of this study bear some potentially far-reaching implications for the use of intermediate-t tracers in paleoceanography: First, whether changes in deep water circulation are recorded at all by radiogenic tracers depends on a sensitive balance between the residence time of the tracer and the water replenishment time of the basin. Second, isotopic compositions in seawater will not always represent their continental source in an unmodified composition. Third, the possibility exists that climate-driven changes in mechanical versus chemical erosion rate result in input of isotope tracers of compositions that vary on the timescales of climate cycles. Such signals can be dispersed basin-wide by rapid lateral advection [Igel and von Blanckenburg, 1999; von Blanckenburg and Igel, 1999] and might obscure isotope variations that result from changes in the thermohaline circulation. It will be important to distinguish between these two causes in paleoceanographic studies.

This conclusion does not invalidate the possible link between the closure of the Panama Gateway and an increase in Labrador Seawater convection. However, it does suggest that in this case, radiogenic isotope tracers would not necessarily record such a change in ocean circulation but are more likely to record changes in weathering associated with the onset of Northern Hemisphere glaciation and increased availability of glacial detritus. This is entirely compatible with observations from longer-t ocean tracers [Hodell et al., 1990; Oxburgh, 1998; Peucker-Ehrenbrink and Blum, 1998].

Acknowledgments. The authors are grateful to the ODP program, in particular the crew and scientific staff of Leg 105, and also Stephen Moorbath for providing samples. Jan Kramers (Berne) is thanked for establishment of excellent new mass spectrometric facilities, Borgi Hebeisen is thanked for technical support, Ben Reynolds is thanked for a contribution to Figure 1, and Benedict De Silva is thanked for his grammatical view of oceanography. The thoughtful unofficial reviews by Martin Frank, Jan Kramers, and Thomas Stocker and constructive official reviews by Bernhard Peucker-Ehrenbrink, Clark Johnson, and an anonymous reviewer improved earlier versions of this paper.

Figure 6. Nd isotope ratios, ¹⁴⁷Sm/¹⁴⁴Nd, and percentage of Nd released during leacheate of dissolved bulk samples (solid symbols) and 0.05 M HCl leacheates (open symbols) of river sediments (circles) and Ocean Drilling Program 645 sediments (squares). The leached fraction represents the Nd available for release into seawater. Analytical details are given in Table 2.

References

- Abouchami, W., S.J.G. Galer, and A. Koschinsky, Pb and Nd isotopes NE Atlantic Fe-Mn crusts: Proxies for paleosources and paleocean circulation, *Geochim. Cosmochim. Acta*, 63, 1489-1505, 1999.
- Albarède, F., A. Simonetti, J.D. Vervoort, J Blichert-Toft, and W. Abouchamie, A Hf-Nd isotopic correlation in ferromanganese nodules. *Geophys. Res. Lett.*, 25, 3895-3898, 1998.
- Andersson, P.S, R. Dahlqvist, J. Ingri and Ö. Gustafsson, The isotopic composition of Nd in a boreal river: A reflection of selective weath-

ering and colloidal transport, Geochim. Cosmochim. Acta, 65, 521-527, 2001. Belshaw, N., P. Freedman, R. O'Nions, M. Frank,

- Belshaw, N., P. Freedman, R. O'Nions, M. Frank, and Y. Guo, A new variable dispersion doublefocusing plasma mass spectrometer with performance illustrated for Pb isotopes, *Int. J. Mass Spectrom.*, 181, 51-58, 1999
- Burton, K.W., H.F. Ling and R.K. O'Nions, Closure of the Central American Isthmus and its effect on deep-water formation in the North Atlantic, *Nature*, 386, 382-385, 1997.
- Burton, K., D. Lee, J. Christensen, A. Halliday,

and J. Hein, Actual timing of neodymium isotopic variations recorded by Fe-Mn crusts in the western North Atlantic, *Earth Planet. Sci. Lett.*, 171, 149-156, 1999.

- Christensen, J.N., A.N. Halliday, L.V. Godfrey, J.R. Hein and D.K. Rea, Climate and ocean dynamics and the lead isotopic records in Pacific ferromanganese crusts, *Science*, 277, 913-918, 1997.
- Cochran, J.K., et al., ²¹⁰Pb scavenging in the North Atlantic and North Pacific Oceans, *Earth Planet. Sci. Lett.*, 97, 332-352, 1990.

i

- Dickson, R.R., and J. Brown, The production of North Atlantic Deep Water: Sources, rates, and pathways, J. Geophys. Res., 99, 12,319-12,341, 1994.
- Erel, Y., Y. Harlavan, and J D. Blum, Lead isotope systematics of granitoid weathering, *Geochim Cosmochim. Acta*, 58, 5299-5306, 1994.
- Frank, M., B.C. Reynolds, and R.K. O'Nions, Nd and Pb isotopes in Atlantic and Pacific water masses before and after closure of the Panama Gateway, *Geology*, 27, 1147-1150, 1999
- Harlavan, Y., Y. Erel, and J.D. Blum, Systematic changes in lead isotopic composition with soil age in glacial granitic terrains, *Geochim Cos*mochim. Acta, 62, 33-46, 1998.
- Haug, G.H., and R. Tiedemann, Effect of the formation of the Isthmus of Panama on Atlantic Ocean thermohaline circulation, *Nature*, 393, 673-676, 1998.
- Heinze, C., and T.J Crowley, Sedimentary response to ocean gateway circulation changes, *Paleoceanography*, 12, 742-745, 1997.
- Hodell, D., G. Mead, and P. Mueller, Variation in the strontium isotopic composition of seawater (8 Ma to present) - Implications for chemicalweathering rates and dissolved fluxes to the oceans, Chem. Geol., 80, 291-307, 1990.
- Igel, H. and F. von Blanckenburg, Lateral mixing and advection of reactive isotope tracers in ocean basins: Numerical modeling, Geochem. Geophys. Geosyst., 1, Paper number 1999GC000003, [10,402 words, 15 figures, 3 tables, 2 movies]. December 13, 1999.
- Kusakabe, M., T.L. Ku, J.R. Southon, and C.I. Measures, Beryllium isotopes in the ocean, *Geochem. J.*, 24, 263-272, 1990.
- Larsen, H.C., A.D. Saunders, P.D. Clift, J. Beget, W. Wei, S. Spezzaferri, and ODP Leg 152 Scientific Party, Seven million years of glaciation in Greenland, *Science*, 264, 952-955, 1994.
- Lee, D.C., A.N. Halliday, J.R. Hen, K.W Burton, J.N. Christensen, and D. Günther, Hafnium isotope stratigraphy of ferromanganese crusts, *Science*, 285, 1052-1054, 1999.
- Mikolajewicz, U. and T.J. Crowley, Response of a coupled ocean/energy balance model to restructed flow through the central American isthmus, *Paleoceanography*, 12, 429-441, 1997.
- Moorbath, S, M. Whitehouse, and B. Kamber, Extreme Nd-isotope heterogeneity in the early

Archean - Fact or fiction? Case studies from northern Canada and west Greenland, *Chem. Geol*, 135, 213-231, 1997.

- Öhlander, B, J. Ingri, M. Land, and H. Schöberg, Change of Sm-Nd isotope composition during weathering of till, *Geochim. Cosmochim. Acta*, 64, 813-820, 2000.
- O'Nions, R.K., M. Frank, F. von Blanckenburg, and H.F. Ling, Secular variations of Nd and Pb isotopes in ferromanganese crusts from the Atlantic, Indian, and Pacific Oceans, Earth Planet. Sci. Lett., 155, 15-28, 1998.
- Oxburgh, R., Variations in the osmium isotope composition of sea water over the past 200,000 years, *Earth Planet. Sci. Lett.*, 159, 183-191, 1998.
- Peucker-Ehrenbrink, B. and J.D. Blum, Re-Os systematics and weathering of Precambrian crustal rocks: Implications for the marine osmium isotope record, *Geochim. Cosmochim. Acta*, 62, 3193-3203, 1998.
- Piepgras, D.J. and G J. Wasserburg, Rare earth element transport in the western North Atlantic inferred from Nd isotopic observations, *Geochim. Cosmochim. Acta*, 51, 1257-1271, 1987.
- Piotrowski, A.M., et al., Changes in erosion and ocean circulation recorded in the Hf isotopic compositions of North Atlantic and Indian Ocean ferromanganese crusts, *Earth Planet. Sci. Lett.*, 181, 315-325, 2000.
- Reynolds, B.C., M. Frank, and R.K. O'Nions, Ndand Pb-isotope time series from Atlantic ferromanganese crusts: Implications for changes in provenance and paleocirculation over the last 8 Myr, Earth Planet. Sci. Lett., 173, 381-396, 1999.
- Rutberg, R.L., S.R. Hemming, and S.L. Goldstein, Reduced North Atlantic Deep Water flux to the glacial Southern Ocean inferred from neodymium isotope ratios, *Nature*, 405, 935-938, 2000.
- Schmitz, W.J., Jr., and M.S. McCartney, On the North Atlantic circulation, *Rev. Geophys.*, 31, 29-49, 1993.
- Silver, L.T., J.A. Woodhead, I.S. Williams, and B.W. Chappell, Uranium in granites from the southwestern United States: Actinide parentdaughter systems, sites and mobilization, *DE*-*ACI3-76GI01664*, Dep. of Energy, Washington D C., 1984.

- Srivastava, S., et al (Ed.), Proceedings of the Ocean Drilling Program, Scientific Results Leg 105, Ocean Drill. Program, College Station, Tex., 1989.
- Stanley, S.M., New horizons for paleontology, with two examples: The rise and fall of the Cretaceous Supertethys and the cause of the modern ice age, J. Paleontol, 69, 999-1007, 1995
- Stordal, M.C. and G.J. Wasserburg, Neodymium isotopic study of Baffin Bay Water: Sources of REE from very old terranes, *Earth Planet. Sci. Lett.*, 77, 259-272, 1986.
- Vance, D, and K. Burton, Neodymium isotopes in planctonic foraminifera: a record of the response of continental weathering and ocean circulation rates to climate change, *Earth Planet. Sci. Lett.*, 173, 365-379, 1999.
- von Blanckenburg, F., Tracing past ocean circulation? Science, 286, 1862-1863, 1999.
- von Blanckenburg, F., and H. Igel, Lateral mixing and advection of reactive isotope tracers in ocean basins: Observations and mechanisms, *Earth Planet. Sci. Lett.*, 169, 113-128, 1999.
- von Blanckenburg, F., and R.K. O'Nions, Response of beryllium and radiogenic isotope ratios in Northern Atlantic Deep Water to the onset of Northern Hemisphere glaciation, Earth Planet. Sci. Lett., 167, 175-182, 1999.
- von Blanckenburg, F, R.K. O'Nions, and J.R. Hein, Distribution and sources of preanthropogenic lead isotopes in deep ocean water from Fe-Mn crusts, *Geochim. Cosmochim. Acta*, 60, 4957-4963, 1996.
- Winter, B.L., C.M. Johnson, and D.L. Clark, Stronthum, neodymum, and lead isotope variations of authigenic and silicate sediment components from the Late Cenozoic Arctic Ocean: Implications for sediment provenance and the source of trace metals in seawater, Geochim. Cosmochim. Acta, 61, 4181-4200, 1997.

T.F. Nägler and F. von Blanckenburg, Isotopengeologie, Mineralogisch-Petrographisches Institut, Universität Bern, Erlachstrasse 9a. CH-3012 Bern, Switzerland. (fvb@mpi.unibe.ch)

(Received June 7, 2000; revised January 20, 2001; accepted March 7, 2001.)