Clinical questions

- 1. Which instrument can be recommended to assess and observe symptoms and symptom related problems?
- 2. At what time and how often should the assessment take place?
- 3. Which aspects should be considered when recommending patients an outcome instrument?
- 4. Which interventions can be recommended to patients concerning handling of surgical wounds?
- 5. What care should be taken after wound healing is complete?
- 6. At what time should consultation take place regarding surgical wound and vulva care?
- 7. Which interventions can be recommended regarding self-management of acute postsurgical pain?
- 8. At what time should consultation take place regarding interventions for acute postsurgical pain?
- 9. Which interventions can be recommended regarding self-management of postsurgical tiredness?
- 10. At what time should consultation take place regarding interventions for postsurgical tiredness?
- 11. Which interventions can be recommended regarding postsurgical urinary symptoms?
- 12. At what time should consultation take place regarding interventions for postsurgical urinary symptoms?
- 13. Which interventions can be recommended regarding self-management of lymphedema?
- 14. At what time should consultation take place regarding interventions for lymphedema?
- 15. Which interventions can be recommended regarding difficulties with sitting?
- 16. Which interventions can be recommended when difficulties occur like vulvar skin irritation, dressing or missing tissue when wearing clothes?
- 17. Which interventions can be recommended regarding difficulties with activities of daily living?
- 18. At what time should a consultation take place regarding difficulties concerning sitting, wearing clothes and activities of daily living?
- 19. Which interventions can be recommended regarding self-management of uncertainty?
- 20. At what time should a consultation take place regarding interventions for uncertainty?
- 21. Which interventions can be recommended regarding coping with changed body and sexuality?
- 22. At what time should a consultation take place regarding interventions for coping with changed body and sexuality?

[table 1: Clinical questions]