

3. Marquard K, Westphal LM, Milki AA, Lathi RB. Etiology of recurrent pregnancy loss in women over the age of 35 years. *Fertil Steril* 2010;**94**: 1473–1477.
4. Sugiura-Ogasawara M, Ozaki Y, Katano K, Suzumori N, Kitaori T, Mizutani E. Abnormal embryonic karyotype is the most frequent cause of recurrent miscarriage. *Hum Reprod* 2012;**27**:2297–2303.

Gayathree Murugappan\* and Ruth Bunker Lathi  
 Division of Reproductive Endocrinology and Infertility, Stanford University, Stanford, CA, USA

\*Correspondence address. E-mail: gayathreem@gmail.com

doi:10.1093/humrep/dew314  
 Advanced Access Publication on December 13, 2016

## Should we also work on an international informed consent for endometriosis surgery?

Sir,

We read the consensus opinion on the recording of deep endometriosis findings at surgery (Vanhie *et al.*, 2016) with great interest. The consensus established by the group of experts will be, without a doubt, extremely useful for comparing surgical techniques and practices. It should have a direct impact on the quality of studies, and should help, as mentioned in the paper, to turn opinion into science. However, I have some concerns on the consensus regarding information provided to the patient.

We believed that, as in many fields in surgery, there should be an internationally agreed informed consent form. This form when agreed upon should be translated into all relevant languages, and provided to international societies, and individual physicians. Consequently, all patients will receive the same information, before surgery about the complications, alternatives, chance of success, risk of recurrence and potential fertility outcomes. An agreed and established international consent form will also be helpful if there is litigation after surgery with a high risk of complications.

Additionally, an identical international informed consent form may improve pre-operative evaluation. Indeed, before signing this form, physicians will have to check if appropriate pre-operative evaluation has been carried out for each potential lesion. Pre-operative evaluation is a key step leading to the success of endometriosis surgery. This informed consent form could be a useful tool to evaluate the efficacy of the pre-operative evaluation and subsequent surgical outcome. Moreover, it should ensure that patients receive accurate and agreed information regarding the surgical procedure.

## Reference

- Vanhie A, Meuleman C, Tomassetti C, Timmerman D, D'Hoore A, Wolthuis A, Van Cleynenbreugel B, Dancet E, Van den Broeck U, Tsalts J *et al.* Consensus on recording deep endometriosis surgery: the CORDES statement. *Hum Reprod* 2016;**31**:1219–1223.

David Soriano and Bouaziz Jerome\*  
 Department of Obstetrics and Gynecology, Sheba Medical Center and Tel-Aviv University-Saklar Medical School, Tel-Aviv, Israel.

\*Correspondence address. Tel: +972-526666469;  
 E-mail: dr.jeromebouaziz@gmail.com

doi:10.1093/humrep/dew335  
 Advanced Access Publication on January 1, 2017

## Reply: Should we also work on an international informed consent for endometriosis surgery?

Sir,

We read with great interest the letter to the editor by Dr David Soriano concerning our recently published article 'Consensus on recording deep endometriosis surgery: the CORDES statement', Vanhie *et al.* (2016). In his letter, Dr Soriano highlights the need for a consensus on an international consent form for endometriosis surgery.

The main goal of the CORDES statement was to provide an instrument for standardized recording of all relevant aspects of deep endometriosis in surgical trials. Although the development of an international informed consent was not within the scope of our project, all co-authors agree that an international consent form for endometriosis surgery would be very useful in daily clinical practice and might lead to an improvement of the preoperative workup.

The systematic review of the literature concerning deep endometriosis, which formed the basis of the CORDES papers, showed that there is an enormous variation in published data on complication rates, success rates and recurrence rates in deep endometriosis surgery. This is due to the lack of standardized definitions, inadequate reporting and the very diverse surgical techniques used. Clearly, results related to clinical outcome after surgery are largely dependent on the population studied, e.g. primary intervention or secondary intervention, and this patient population is often inadequately characterized. In our view, the lack of good information from high quality trials impedes the development of an international consent form at present.

Based on our experience, the development of an internationally accepted consent form, applicable in a wide range of countries, will require a long and difficult consensus process. As stated earlier, this was not the ambition of the authors publishing the CORDES statement, but may be a next step for these authors, or for other groups like the World Endometriosis Society, European Endometriosis League and/or Society for Endometriosis and Endometrial Disorders.

## Reference

- Vanhie A, Meuleman C, Tomassetti C, Timmerman D, D'Hoore A, Wolthuis A, Van Cleynenbreugel B, Dancet E, Van den Broeck U, Tsalts J, Renner SP, Ebert AD, Carmona F, Abbott J, Stepniowska A, Taylor H, Saridogan E, Mueller M, Keckstein J, Pluchino N, Janik G, Zupi E, Minelli L, Cooper M, Dunselman G, Koh C, Abrao M, Chapron C, D'Hooghe T. Consensus on recording deep endometriosis surgery: the CORDES statement. *Hum Reprod* 2016 Apr 19. pii: dew067.

A. Vanhie<sup>1,\*</sup>, C. Meuleman<sup>1</sup>, C. Tomassetti<sup>1</sup>, D. Timmerman<sup>1</sup>, A. D'Hoore<sup>2</sup>,  
 A. Wolthuis<sup>2</sup>, B. Van Cleynenbreugel<sup>3</sup>, E. Dancet<sup>1</sup>, U. Van den Broeck<sup>1</sup>,  
 J. Tsaltas<sup>4</sup>, S.P. Renner<sup>5</sup>, A.D. Ebert<sup>6</sup>, F. Carmona<sup>7</sup>, J. Abbott<sup>8</sup>,  
 A. Stepniewska<sup>9</sup>, H. Taylor<sup>10</sup>, E. Saridogan<sup>11</sup>, M. Mueller<sup>12</sup>, J. Keckstein<sup>13</sup>,  
 N. Pluchino<sup>14</sup>, E. Zupi<sup>15</sup>, G. Dunselman<sup>16,17</sup>, M.S. Abrao<sup>18,19</sup>, C. Chapron<sup>20,21</sup>,  
 and T. D'Hooghe<sup>1,22,23</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, University Hospital Leuven, Belgium

<sup>2</sup>Department of Abdominal Surgery, University Hospital Leuven, Belgium

<sup>3</sup>Department of Urology, University Hospital Leuven, Belgium

<sup>4</sup>Monash Health and Monash University, Melbourne, Victoria, Australia

<sup>5</sup>Frauenklinik, Universitätsklinikum Erlangen, Germany

<sup>6</sup>Praxis für Frauengesundheit, Gynäkologie und Geburtshilfe, Berlin, Germany

<sup>7</sup>Service of Gynecology, Hospital Clinic of Barcelona, University of Barcelona, Spain

<sup>8</sup>Royal Hospital for Women and University of New South Wales, Sydney, Australia

<sup>9</sup>Department of Obstetrics and Gynecology, Ospedale Sacro Cuore, Verona, Italy

<sup>10</sup>Yale School of Medicine, New Haven, CT, USA

<sup>11</sup>University College London Hospitals, London, UK

<sup>12</sup>Universitätsklinik für Frauenheilkunde, Universitätsspital Bern, Switzerland

<sup>13</sup>Landeskrankenanstalten-Betriebsgesellschaft (KABEG) and Landeskrankenhaus  
 Villach, Abteilung für Gynäkologie und Geburtshilfe, Villach, Austria

<sup>14</sup>Department of Obstetrics and Gynecology, University Hospital of Geneva,  
 Switzerland

<sup>15</sup>University of Siena, Siena, Italy

<sup>16</sup>Department of Obstetrics and Gynaecology, Research Institute GROW, Maastricht,  
 The Netherlands

<sup>17</sup>University Medical Centre, PO Box 5800, 6202 AZ Maastricht, The Netherlands

<sup>18</sup>Division of Reproductive Medicine of Sirio Libanes Hospital, Sao Paulo, Brazil

<sup>19</sup>Division of Endometriosis, Department of Obstetrics and Gynaecology São Paulo  
 University, São Paulo, Brazil

<sup>20</sup>Faculty of Medicine Université Paris Descartes, Sorbonne Paris Cité, Paris, France

<sup>21</sup>Department of Gynecology, Obstetrics, and Reproductive Medicine, Centre  
 Hospitalier Universitaire Cochin of the Groupe Hospitalier Universitaire Ouest, Paris,  
 France

<sup>22</sup>Faculty of Medicine, Leuven University, Belgium

<sup>23</sup>Faculty of Medicine, Yale University, New Haven, CT, USA

\*Correspondence address. Arne Vanhie. E-mail: arnevanhie@hotmail.com

doi:10.1093/humrep/dew336

Advanced Access Publication on January 1, 2017